

The University of Bradford Institutional Repository

<http://bradscholars.brad.ac.uk>

This work is made available online in accordance with publisher policies. Please refer to the repository record for this item and our Policy Document available from the repository home page for further information.

To see the final version of this work please visit the publisher's website. Access to the published online version may require a subscription.

Link to publisher's version: <http://dx.doi.org/10.1016/j.bpj.2013.11.888>

Citation: Murali K, Bollepalli MK, Fowler PW et al (2014) Structural and Thermodynamic Characterization of the Gating Pathway in a K⁺ Channel. *Biophysical Journal*. 106(2) Supplement 1: 155A.

Copyright statement: © 2014 Elsevier. Reproduced in accordance with the publisher's self-archiving policy. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#).



Structural and Thermodynamic Characterization of the Gating Pathway in a K^D Channel

Murali K. Bollepalli¹, Philip W. Fowler², Markus Rapedius¹, Lijun Shang³, Mark S.P. Sansom², Stephen J. Tucker³, Thomas Baukrowitz¹.

¹Institute of Physiology, University of Kiel, Germany,

²Department of Biochemistry, University of Oxford, United Kingdom,

³Clarendon Laboratory, Department of Physics, University of Oxford, United Kingdom.

Structures of inwardly-rectifying (Kir) potassium channels are now available in many different crystallographic states. We now analyse these structures in the context of functional data for mutations at over 180 positions within the Kir1.1 (ROMK) channel. This reveals an extensive network of physically interacting residues which stabilise the pre-open and open-states of the channel, but which breaks down upon channel closure. This approach not only validates a structural gating pathway for the Kir channel, but also provides insight into the structure of the transition state connecting these crystallographic states. This gating network also appears to be an important structural determinant of the thermodynamic stability of these different gating states as well as influencing the impact of mutations on channel function, and suggest that such state-dependent physical connectivity between residues may also be relevant to our understanding of other allosteric proteins.