

(Adelson & Bergen, 1985, Anstis, 1970, Attneave, 1974, Bair, Zohary & Newsome, 2001, Baker & Braddick, 1982, Baker & Braddick, 1985, Baker, Hess & Zihl, 1991, Baker, Petersen, Newsome & Allman, 1981, Barlow & Tripathy, 1997, Bischof & Dillolo, 1990, Bisley, Zaksas, Droll & Pasternak, 2004, Blasdel & Fitzpatrick, 1984, Born & Bradley, 2005, Braddick, 1974, Braddick, 1980, Braddick, O'Brien, Wattam-Bell, Atkinson, Hartley & Turner, 2001, Braddick, Ruddock, Morgan & Marr, 1980, Britten, Shadlen, Newsome & Movshon, 1992, Brown & He, 2000, Burgess, 1990, Burr, 1979, Cavanagh, Boeglin & Favreau, 1985, Cavanagh & Mather, 1989, Chang & Julesz, 1983, Cleary & Braddick, 1990a, Cleary & Braddick, 1990b, Devalois, Yund & Hepler, 1982, Douglas, Neve, Quittenbaum, Alam & Prusky, 2006, Downing & Movshon, 1989, Dubner & Zeki, 1971, Eagle, 1996, Eagle & Rogers, 1996, Eagle & Rogers, 1997, Edwards, Badcock & Nishida, 1996, Exner, 1888, Fitzpatrick, Lund & Blasdel, 1985, Foster, Thorson, McIlwain & Biedermanthorson, 1981, Fredericksen, Verstraten & Vandegrind, 1994, Girard, Salin & Bullier, 1992, Goodale & Milner, 1992, Graham, Baker, Hetch & Lloyd, 1948, Groh, 2000, Hadani, Gur, Meiri & Fender, 1980, Hawken, Parker & Lund, 1988, Hess, Baker & Zihl, 1989, Howell & Hess, 1978, Hubel & Wiesel, 1959, Hubel & Wiesel, 1962, Hubel & Wiesel, 1968, Hubel & Wiesel, 1972)(Julesz, 1960, Julesz & Payne, 1968, Klam, Zemel & Pouget, 2008, Kolers, 1972, Korte, 1915, Kulikowski & Vidyasagar, 1984, Lappin & Bell, 1976, Lu & Sperling, 1995, Marr & Ullman, 1981, Maunsell & Van Essen, 1983, Merigan & Maunsell, 1993, Mikami, 1992, Morgan, 1992, Morgan & Fahle, 1992, Morgan, Perry & Fahle, 1997, Movshon, Adelson, Gizzi & Newsome, 1985, Movshon & Newsome, 1996, Nakayama, 1985, Nakayama & Silverman, 1984, Nakayama & Tyler, 1981, Nassi & Callaway, 2005, Newsome & Pare, 1988, Nishida & Sato, 1991, Orban, Kennedy & Bullier, 1986, Pelli & Farell, 1999, Priebe, Lisberger & Movshon, 2006, Purushothaman & Bradley, 2005, Ramachandran & Anstis, 1983, Reichardt, 1961, Rodman, Gross & Albright, 1989)(Salmon, 1970, Sato, 1990, Scase, Braddick & Raymond, 1996, Schiller, Finlay & Volman, 1976, Sekuler, 1996, Shadlen & Carney, 1986, Smith & Ledgeway, 2001, Stratton, 1902, Todd & Norman, 1995, Treue, Hol & Rauber, 2000, Tripathy, 2002, Tripathy & Barlow, 2001, Ullman, 1979, Ungerleider & Mishkin, 1982, van de Grind, Koenderink & van Doorn, 1986, van de Grind, Konderink & van Doorn, 1987, van de Grind, van Doorn & Koenderink, 1983, van Doorn & Koenderink, 1982, van Doorn & Koenderink, 1984, van Essen, Maunsell & Bixby, 1981, van Santen & Sperling, 1985, von Grunau, 1986, Watson & Ahumada, 1985, Wertheimer, 1912, Wertheimer, 1961, Westheimer, 1979, Zaksas & Pasternak, 2005, Zeki, 1974, Zihl, von Cramon & Mai, 1983, Zohary, Shadlen & Newsome, 1994)

References

- Adelson, E.H., & Bergen, J.R. (1985). Spatiotemporal energy models for the perception of motion. *J Opt Soc Am A*, 2 (2), 284-299.
- Anstis, S.M. (1970). Phi movement as a subtraction process. *Vision Research*, 10 (12), 1411-1430.
- Attneave, F. (1974). Apparent movement and what-where connection. *Psychologia*, 17 (2), 108-120.
- Bair, W., Zohary, E., & Newsome, W.T. (2001). Correlated firing in macaque visual area MT: Time scales and relationship to behaviour. *Journal of Neuroscience*, 21 (5), 1676-1697.
- Baker, C.L., & Braddick, O.J. (1982). The basis of area and dot number effects in random dot motion perception. *Vision Research*, 22 (10), 1253-1259.
- Baker, C.L., & Braddick, O.J. (1985). Eccentricity-dependent scaling of the limits for short-range apparent motion perception. *Vision Research*, 25 (6), 803-812.
- Baker, C.L., Jr., Hess, R.F., & Zihl, J. (1991). Residual motion perception in a "motion-blind" patient, assessed with limited-lifetime random dot stimuli. *J Neurosci*, 11 (2), 454-461.
- Baker, J.F., Petersen, S.E., Newsome, W.T., & Allman, J.M. (1981). Visual response properties of neurons in 4 extrastriate visual areas of the owl monkey (*Aotus-trivirgatus*) - a quantitative comparison of medial, dorsomedial, dorsolateral, and middle temporal areas. *Journal of Neurophysiology*, 45 (3), 397-416.
- Barlow, H., & Tripathy, S.P. (1997). Correspondence noise and signal pooling in the detection of coherent visual motion. *Journal of Neuroscience*, 17 (20), 7954-7966.
- Bischof, W.F., & Dilollo, V. (1990). Perception of directional sampled motion in relation to displacement and spatial-frequency - evidence for a unitary motion system. *Vision Research*, 30 (9), 1341-1362.

- Bisley, J.W., Zaksas, D., Droll, J.A., & Pasternak, T. (2004). Activity of neurons in cortical area MT during a memory for motion task. *Journal of Neurophysiology*, *91* (1), 286-300.
- Blasdel, G.G., & Fitzpatrick, D. (1984). Physiological organization of layer-4 in macaque striate cortex. *Journal of Neuroscience*, *4* (3), 880-895.
- Born, R.T., & Bradley, D.C. (2005). Structure and function of visual area MT. *Annu Rev Neurosci*, *28*, 157-189.
- Braddick, O. (1974). Short-range process in apparent motion. *Vision Research*, *14* (7), 519-527.
- Braddick, O.J. (1980). Low-level and high-level processes in apparent motion. *Philosophical Transactions of the Royal Society of London B*, *290*, 137-151.
- Braddick, O.J., O'Brien, J.M.D., Wattam-Bell, J., Atkinson, J., Hartley, T., & Turner, R. (2001). Brain areas sensitive to coherent visual motion. *Perception*, *30* (1), 61-72.
- Britten, K.H., Shadlen, M.N., Newsome, W.T., & Movshon, J.A. (1992). The analysis of visual-motion - a comparison of neuronal and psychophysical performance. *Journal of Neuroscience*, *12* (12), 4745-4765.
- Brown, R.O., & He, S. (2000). Visual motion of missing-fundamental patterns: motion energy versus feature correspondence. *Vision Research*, *40* (16), 2135-2147.
- Burgess, A.E. (1990). High level visual decision efficiencies. In: C. Blakemore (Ed.) *Vision: Coding and efficiency* (pp. 431-440). New York: Cambridge University Press.
- Burr, D.C. (1979). Acuity for apparent vernier offset. *Vision Research*, *19* (7), 835-837.
- Cavanagh, P., Boeglin, J., & Favreau, O.E. (1985). Perception of motion in equiluminous kinematograms. *Perception*, *14* (2), 151-162.
- Cavanagh, P., & Mather, G. (1989). Motion: The long and the short of it. *Spatial Vision*, *4*, 103-129.
- Chang, J.J., & Julesz, B. (1983). Displacement limits for spatial frequency filtered random-dot cinematograms in apparent motion. *Vision Research*, *23*, 1379-1385.

- Cleary, R., & Braddick, O.J. (1990a). Direction discrimination for band pass filtered random dot kinematograms. *Vision Research*, 30, 303-316.
- Cleary, R., & Braddick, O.J. (1990b). Masking of low-frequency information in short-range apparent motion. *Vision Research*, 30 (2), 317-327.
- Devalois, R.L., Yund, E.W., & Hepler, N. (1982). The orientation and direction selectivity of cells in macaque visual-cortex. *Vision Research*, 22 (5), 531-544.
- Douglas, R.M., Neve, A., Quittenbaum, J.P., Alam, N.M., & Prusky, G.T. (2006). Perception of visual motion coherence by rats and mice. *Vision Research*, 46 (18), 2842-2847.
- Downing, C.J., & Movshon, J.A. (1989). Spatial and temporal summation in the detection of motion in stochastic random dot displays. *Invest Ophthalmol and Visual Sci (Suppl)*, 30, 72.
- Dubner, R., & Zeki, S.M. (1971). Response properties and receptive fields of cells in an anatomically defined region of the superior temporal sulcus in the monkey. *Brain Res*, 35 (2), 528-532.
- Eagle, R.A. (1996). What determines the maximum displacement limit for spatially broadband kinematograms? *Journal of the Optical Society of America A-Optics Image Science and Vision*, 13 (3), 408-418.
- Eagle, R.A., & Rogers, B.J. (1996). Motion detection is limited by element density not spatial frequency. *Vision Research*, 36 (4), 545-558.
- Eagle, R.A., & Rogers, B.J. (1997). Effects of dot density, patch size and contrast on the upper spatial limit for direction discrimination in random-dot kinematograms. *Vision Research*, 37 (15), 2091-2102.
- Edwards, M., Badcock, D.R., & Nishida, S. (1996). Contrast sensitivity of the motion system. *Vision Research*, 36 (16), 2411-2421.
- Exner, S. (1888). Uber optische bewegungsempfindungen. *Biologische Centralblatt*, 8, 437-448.

- Fitzpatrick, D., Lund, J.S., & Blasdel, G.G. (1985). Intrinsic connections of macaque striate cortex: afferent and efferent connections of lamina 4C. *J Neurosci*, 5 (12), 3329-3349.
- Foster, D.H., Thorson, J., McIlwain, J.T., & Biedermanthorson, M. (1981). The fine-grain movement illusion - a perceptual probe of neuronal connectivity in the human visual-system. *Vision Research*, 21 (7), 1123-1128.
- Fredericksen, R.E., Verstraten, F.A.J., & Vandegrind, W.A. (1994). Spatial summation and its interaction with the temporal integration mechanism in human motion perception. *Vision Research*, 34 (23), 3171-3188.
- Girard, P., Salin, P.A., & Bullier, J. (1992). Response selectivity of neurons in area-MT of the macaque monkey during reversible inactivation of area-V1. *Journal of Neurophysiology*, 67 (6), 1437-1446.
- Goodale, M.A., & Milner, A.D. (1992). Separate visual pathways for perception and action. *Trends Neurosci*, 15 (1), 20-25.
- Graham, C.H., Baker, K.E., Hetch, M., & Lloyd, V.V. (1948). Factors influencing thresholds for monocular movement parallax. *Journal of Experimental Psychology*, 38 (3), 205-223.
- Groh, J.M. (2000). Predicting perception from population codes. *Nature Neuroscience*, 3 (3), 201-202.
- Hadani, I., Gur, M., Meiri, A.Z., & Fender, D.H. (1980). Hyperacuity in the detection of absolute and differential displacements of random dot patterns. *Vision Research*, 20 (11), 947-951.
- Hawken, M.J., Parker, A.J., & Lund, J.S. (1988). Laminar organization and contrast sensitivity of direction-selective cells in the striate cortex of the Old World monkey. *J Neurosci*, 8 (10), 3541-3548.
- Hess, R.H., Baker, C.L., Jr., & Zihl, J. (1989). The "motion-blind" patient: low-level spatial and temporal filters. *J Neurosci*, 9 (5), 1628-1640.

- Howell, E.R., & Hess, R.F. (1978). Functional area for summation to threshold for sinusoidal gratings. *Vision Research*, 18 (4), 369-374.
- Hubel, D.H., & Wiesel, T.N. (1959). Receptive fields of single neurones in the cat's striate cortex. *Journal of Physiology*, 148, 574-591.
- Hubel, D.H., & Wiesel, T.N. (1962). Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *Journal of Physiology*, 160, 106-165.
- Hubel, D.H., & Wiesel, T.N. (1968). Receptive fields and functional architecture of monkey striate cortex. *Journal of Physiology (London)*, 195, 215-243.
- Hubel, D.H., & Wiesel, T.N. (1972). Laminar and columnar distribution of geniculo-cortical fibers in the macaque monkey. *J Comp Neurol*, 146 (4), 421-450.
- Julesz, B. (1960). Binocular depth perception of computer-generated patterns. *Bell System Tech J*, 39, 1125-1162.
- Julesz, B., & Payne, R.A. (1968). Differences between monocular and binocular stroboscopic movement perception. *Vision Research*, 8, 433-444.
- Klam, F., Zemel, R.S., & Pouget, A. (2008). Population coding with motion energy filters: The impact of correlations. *Neural Computation*, 20 (1), 146-175.
- Kolers, P.A. (1972). Aspects of motion perception. Sydney: Pergamon Press.
- Korte, A. (1915). Kinematoskopische untersuchungen [Kinematoscopic investigations] *Zeitschrift für Psychologie*, 72, 194-296.
- Kulikowski, J.J., & Vidvasagar, T.R. (1984). Macaque striate cortex - pattern, movement and color processing. *Ophthalmic and Physiological Optics*, 4 (1), 77-81.
- Lappin, J.S., & Bell, H.H. (1976). Detection of coherence in moving random-dot patterns. *Vision Research*, 16 (2), 161-168.
- Lu, Z.L., & Sperling, G. (1995). The functional architecture of human visual motion perception. *Vision Research*, 35 (19), 2697-2722.

- Marr, D., & Ullman, S. (1981). Directional selectivity and its use in early visual processing. *Proceedings of the Royal Society of London (B)*, 211, 151-180.
- Maunsell, J.H.R., & Van Essen, D.C. (1983). Functional properties of neurons in the middle temporal visual area of the macaque monkey. *Journal of Neurophysiology*, 49, 1127-1146.
- Merigan, W.H., & Maunsell, J.H. (1993). How parallel are the primate visual pathways? *Annu Rev Neurosci*, 16, 369-402.
- Mikami, A. (1992). Spatiotemporal characteristics of direction-selective neurons in the middle temporal visual area of the macaque monkeys. *Experimental Brain Research*, 90 (1), 40-46.
- Morgan, M.J. (1992). Spatial filtering precedes motion detection. *Nature*, 355, 344-346.
- Morgan, M.J., & Fahle, M. (1992). Effects of pattern element density upon displacement limits for motion detection in random binary luminance patterns. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 248 (1322), 189-198.
- Morgan, M.J., Perry, R., & Fahle, M. (1997). The spatial limit for motion detection in noise depends on element size, not on spatial frequency. *Vision Research*, 37 (6), 729-736.
- Movshon, J.A., Adelson, E.H., Gizzi, M.S., & Newsome, W. (1985). The analysis of moving patterns. In: C. Chagas, R. Gattas, & C. Gross (Eds.), *Pattern recognition mechanisms* Berlin: Springer-Verlag.
- Movshon, J.A., & Newsome, W.T. (1996). Visual response properties of striate cortical neurons projecting to area MT in macaque monkeys. *Journal of Neuroscience*, 16 (23), 7733-7741.
- Nakayama, K. (1985). Biological image motion processing: a review. *Vision Research*, 25, 625-660.
- Nakayama, K., & Silverman, G.H. (1984). Temporal and spatial characteristics of the upper displacement limit for motion in random dots. *Vision Research*, 24, 293-299.

- Nakayama, K., & Tyler, C.W. (1981). Psychophysical isolation of movement sensitivity by removal of familiar position cues. *Vision Research*, 21, 427-433.
- Nassi, J.J., & Callaway, E.M. (2005). Indirect parvocellular input to the area MT of macaque monkey. *Society for Neuroscience Abstracts, Washington DC*, Programme number 1377.
- Newsome, W.T., & Pare, E.B. (1988). A selective impairment of motion perception following lesions of the middle temporal visual area (MT). *Journal of Neuroscience*, 8, 2201-2211.
- Nishida, S., & Sato, T. (1991). Positive motion aftereffect induced by a bandpass random dot kinematogram. *Investigative Ophthalmology & Visual Science*, 32 (4), 827-827.
- Orban, G.A., Kennedy, H., & Bullier, J. (1986). Velocity sensitivity and direction selectivity of neurons in area-V1 and area-V2 of the monkey - influence of eccentricity. *Journal of Neurophysiology*, 56 (2), 462-480.
- Pelli, D.G., & Farell, B. (1999). Why use noise? *Journal of the Optical Society of America A-Optics Image Science and Vision*, 16 (3), 647-653.
- Priebe, N.J., Lisberger, S.G., & Movshon, J.A. (2006). Tuning for spatiotemporal frequency and speed in directionally selective neurons of macaque striate cortex. *Journal of Neuroscience*, 26 (11), 2941-2950.
- Purushothaman, G., & Bradley, D.C. (2005). Neural population code for fine perceptual decisions in area MT. *Nature Neuroscience*, 8 (1), 99-106.
- Ramachandran, V.S., & Anstis, S.M. (1983). Displacement thresholds for coherent apparent motion in random dot-patterns. *Vision Research*, 23 (12), 1719-1724.
- Reichardt, W. (1961). Autocorrelation, a principle for the evaluation of sensory information by the central nervous system. In: W.A. Rosenblith (Ed.) *Sensory communication* (pp. 303-317). New York: Wiley.

- Rodman, H.R., Gross, C.G., & Albright, T.D. (1989). Afferent basis of visual response properties in area MT of the macaque .I. effects of striate cortex removal. *Journal of Neuroscience*, 9 (6), 2033-2050.
- Salmon, W.C. (1970). Zeno's paradoxes. New York: Bobbs-Merrill.
- Sato, T. (1990). Effects of dot size and dot density on motion perception with random dot kinematograms. *Perception (Suppl.)*, 19, 329.
- Scase, M.O., Braddick, O.J., & Raymond, J.E. (1996). What is noise for the motion system? *Vision Research*, 36 (16), 2579-2586.
- Schiller, P., Finlay, B.L., & Volman, S.F. (1976). Quantitative studies of single-cell properties in monkey striate cortex. I. Spatiotemporal organization of receptive fields. *Journal of Neurophysiology*, 39, 1288-1319.
- Sekuler, R. (1996). Motion perception: A modern view of Wertheimer's 1912 monograph. *Perception*, 25, 1243-1258.
- Shadlen, M., & Carney, T. (1986). Mechanisms of human motion perception revealed by a new cyclopean illusion. *Science*, 232 (4746), 95-97.
- Smith, A.T., & Ledgeway, T. (2001). Motion detection in human vision: a unifying approach based on energy and features. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 268 (1479), 1889-1899.
- Stratton, C. (1902). Visible motion and space threshold. *Psychology Review*, 9, 433-447.
- Todd, J.T., & Norman, J.F. (1995). The effects of spatiotemporal integration on maximum displacement thresholds for the detection of coherent motion. *Vision Research*, 35 (16), 2287-2302.
- Treue, S., Hol, K., & Rauber, H.J. (2000). Seeing multiple directions of motion - physiology and psychophysics. *Nature Neuroscience*, 3 (3), 270-276.

- Tripathy, S.P. (2002). Correspondence noise and Dmin in random dot kinematograms. *Journal of Vision*, 2 (7), 741a.
- Tripathy, S.P., & Barlow, H.B. (2001). Unified theory for threshold coherence and Dmax in random dot kinematograms. *Perception (Suppl.)*, 30, 32.
- Ullman, S. (1979). The interpretation of visual motion. Cambridge, MA: MIT Press.
- Ungerleider, & Mishkin (1982). Analysis of visual behavior. Cambridge, MA: MIT Press.
- van de Grind, W.A., Koenderink, J.J., & van Doorn, A.J. (1986). The distribution of human motion detector properties in the monocular visual field. *Vision Research*, 26, 797-810.
- van de Grind, W.A., Konderink, J.J., & van Doorn, A.J. (1987). Influence of contrast on foveal and peripheral detection of coherent motion in moving random-dot patterns. *Journal of the Optical Society of America A*, 4, 1643-1652.
- van de Grind, W.A., van Doorn, A.J., & Koenderink, J.J. (1983). Detection of coherent motion in peripherally viewed random dot patterns. *Journal of the Optical Society of America*, 73, 1674-1683.
- van Doorn, A.J., & Koenderink, J.J. (1982). Spatial properties of the visual detectability of moving spatial white noise. *Experimental Brain Research*, 45, 189-195.
- van Doorn, A.J., & Koenderink, J.J. (1984). Spatiotemporal integration in the detection of coherent motion. *Vision Research*, 24, 47-53.
- van Essen, D.C., Maunsell, J.H.R., & Bixby, J.L. (1981). The middle temporal visual area in the macaque - myeloarchitecture, connections, functional-properties and topographic organization. *Journal of Comparative Neurology*, 199 (3), 293-326.
- van Santen, J.P.H., & Sperling, G. (1985). Elaborated Reichardt detectors. *Journal of the Optical Society of America A*, 2, 300-321.
- von Grunau, M.W. (1986). A motion aftereffect for long-range stroboscopic apparent motion. *Perception & Psychophysics*, 40 (1), 31-38.

- Watson, A.B., & Ahumada, A.J. (1985). Model of human visual-motion sensing. *Journal of the Optical Society of America A-Optics Image Science and Vision*, 2 (2), 322-342.
- Wertheimer, M. (1912). Untersuchungen zur lehre der Gestalt. *Psychologische Forschung*, 4, 301-350.
- Wertheimer, M. (1961). Experimentelle studen uber das sehen von bewegung. (Zeitschrift fur Psychologie, 61, 161-265). In: T. Shipley (Ed.) *Classics in psychology* (New York: Philosophical Library).
- Westheimer, G. (1979). Spatial sense of the eye - Proctor Lecture. *Investigative Ophthalmology & Visual Science*, 18 (9), 893-912.
- Zaksas, D., & Pasternak, T. (2005). Area MT neurons respond to visual motion distant from their receptive fields. *Journal of Neurophysiology*, 94 (6), 4156-4167.
- Zeki, S.M. (1974). Functional organization of a visual area in posterior bank of superior temporal sulcus of rhesus-monkey. *Journal of Physiology-London*, 236 (3), 549-73.
- Zihl, J., von Cramon, D., & Mai, N. (1983). Selective disturbance of movement vision after bilateral brain damage. *Brain*, 106 (Pt 2), 313-340.
- Zohary, E., Shadlen, M.N., & Newsome, W.T. (1994). Correlated neuronal discharge rate and its implications for psychophysical performance. *Nature*, 370, 140-143.