

LIST OF TABLES

<i>Table 1. Human keratins old and new designations</i>	16
<i>Table 2. Immunoglobulin functions and concentrations in human serum.</i>	38
<i>Table 3. IgG concentration purified from normal and AA serum by UV spectrophotometry</i>	79
<i>Table 4. Stepped voltage regime used in IEF</i>	86
<i>Table 5. Proteins identified by MALDI-TOF-MS from 2D electrophoresis gels of immunoprecipitate of 62 μL of AA12 serum and 250 μg of HF extract</i>	112
<i>Table 6. List of proteins identified from 2D electrophoresis gel spots of HF extract</i>	121
<i>Table 7. List of proteins identified by LC-MALDI-MS/MS, of immunoprecipitate of HF extract and whole AA5 serum</i>	129
<i>Table 8. Proteins identified by MALDI-TOF/TOF in 20 AA and normal samples and 3 negative controls with Mascot scores over 30</i>	139
<i>Table 9. Proteins identified by MALDI-TOF/TOF with Mascot scores over 30, excluding IgGs, false positives and those that only observed in negative controls</i>	145
<i>Table 10. Frequency of occurrence, mean score, SEM and significance of score difference between the AA and normal groups for 6 proteins identified by MALDI-TOF/TOF</i>	145
<i>Table 11. Amount of guinea pig serum complement, human serum and commercial antibodies used for HF culture experiments</i>	153
<i>Table 12. Blocking sera, primary and secondary antibodies used in double/single immunofluorescence analysis</i>	169
<i>Table 13. Summary of reactivity of AA sera and normal sera antibodies to normal human scalp HF by indirect immunofluorescence – by structure and co-localisation with THH</i>	188