

REFERENCES

- Aderem, A. & Underhill, D. M. (1999). Mechanisms of phagocytosis in macrophages. *Annual Review of Immunology*, **17**, 593-623.
- Aebi, U., Fowler, W. E., Rew, P. & Sun, T. T. (1983). The fibrillar substructure of keratin filaments unraveled. *Journal of Cell Biology*, **97**, 1131-1143.
- Albrethsen, J. (2007). Reproducibility in protein profiling by MALDI-TOF mass spectrometry. *Clinical Chemistry*, **53**, 852-858.
- Andersen-Ranberg, K., Hoier-Madsen, M., Wiik, A., Jeune, B. & Hegedus, L. (2004). High prevalence of autoantibodies among Danish centenarians. *Clinical and Experimental Immunology*, **138**, 158-163.
- Anhalt, G. J., Till, G. O., Diaz, L. A., Labib, R. S., Patel, H. P. & Eaglstein, N. F. (1986). Defining the role of complement in experimental pemphigus vulgaris in mice. *Journal of Immunology*, **137**, 2835-2840.
- Arck, P. C., Handjiski, B., Peters, E. M. J., Peter, A. S., Hagen, E., Fischer, A., Klapp, B. F. & Paus, R. (2003). Stress inhibits hair growth in mice by induction of premature catagen development and deleterious perifollicular inflammatory events via neuropeptide substance P-dependent pathways. *American Journal of Pathology*, **162**, 803-814.
- Arlaud, G. J., Gaboriaud, C., Thielens, N. M., Budayova-Spano, M., Rossi, V. & Fontecilla-Camps, J. C. (2002). Structural biology of the C1 complex of complement unveils the mechanisms of its activation and proteolytic activity. *Molecular Immunology*, **39**, 383-394.
- Arteaga, L. A., Prisayanh, P. S., Warren, S. J. P., Liu, Z., Diaz, L. A. & Lin, M. S. (2002). A subset of pemphigus foliaceus patients exhibits pathogenic autoantibodies against both desmoglein-1 and desmoglein-3. *Journal of Investigative Dermatology*, **118**, 806-811.

- Ashcroft, A. E. (1997). Ionization methods in organic mass spectrometry. Edited by Barnett, N. W. The Royal Society of Chemistry, Cambridge.
- Atassi, M. Z. & Casali, P. (2008). Molecular mechanisms of autoimmunity. *Autoimmunity*, **41**, 123-132.
- Avrameas, S. (1991). Natural autoantibodies: from 'horror autotoxicus' to 'gnothi seauton'. *Immunology Today*, **12**, 154-159.
- Bach, F. H. (1991). The role of genetics in autoimmune disease. *Journal of Investigative Dermatology*, **96**, S86-S86.
- Bagshaw, R. D., Callahan, J. W. & Mahuran, D. J. (2000). Desalting of in-gel-digested protein sample with mini-C18 columns for matrix-assisted laser desorption ionization time of flight peptide mass fingerprinting. *Analytical Biochemistry*, **284**, 432-435.
- Barahmani, N., Whaley, K. & Duvic, M. (2003). Alopecia areata after allogeneic bone marrow transplantation from an affected, human leukocyte antigen-matched sibling. *Journal of the American Academy of Dermatology*, **49**, 1192-1192.
- Barahmani, N., de Andrade, M., Slusser, J. P., Zhang, Q. & Duvic, M. (2006). Major histocompatibility complex class I chain-related gene A polymorphisms and extended haplotypes are associated with familial alopecia areata. *Journal of Investigative Dermatology*, **126**, 74-78.
- Barker, C. F. & Billingham, R. E. (1977). Immunologically privileged sites. *Advances in Immunology*, **25**, 1-54.

- Baumann, H. & Gauldie, J. (1994). The Acute-Phase Response. *Immunology Today*, **15**, 74-80.
- Beardi, B., Wanert, F., Zoller, M., Freyschmidt-Paul, P., Bodemer, W. & Kaup, F. J. (2007). Alopecia areata in a rhesus monkey (*Macaca mulatta*). *Journal of Medical Primatology*, **36**, 124-130.
- Bentley, D. R. (1986). Primary structure of human complement component C2 - Homology to two unrelated protein families. *Biochemical Journal*, **239**, 339-345.
- Bernot, K. M., Coulombe, P. A. & McGowan, K. M. (2002). Keratin 16 expression defines a subset of epithelial cells during skin morphogenesis and the hair cycle. *Journal of Investigative Dermatology*, **119**, 1137-1149.
- Berzofsky, J. A. & Terabe, M. (2008). NKT cells in tumor immunity: Opposing subsets define a new immunoregulatory axis. *Journal of Immunology*, **180**, 3627-3635.
- Betterle, C., Greggio, N. A. & Volpato, M. (1998). Clinical review 93: Autoimmune polyglandular syndrome type 1. *Journal of Clinical Endocrinology & Metabolism*, **83**, 1049-1055.
- Betz, R. C., König, K., Flaquer, A., Redler, S., Eigelshoven, S., Kortum, A. K., Hanneken, S., Hillmer, A., Tuting, T., Lambert, J., De Weert, J., Kruse, R., Lutz, G., Blaumeiser, B. & Nothen, M. M. (2008). The R620W polymorphism in PTPN22 confers general susceptibility for the development of alopecia areata. *British Journal of Dermatology*, **158**, 389-391.

- Birbeck, M. S. C. & Mercer, E. H. (1957). The electron microscopy of the human hair follicle .3. The inner root sheath and trichohyaline. *Journal of Biophysical & Biochemical Cytology*, **3**, 223-230.
- Biron, C. A., Nguyen, K. B., Pien, G. C., Cousens, L. P. & Salazar-Mather, T. P. (1999). Natural killer cells in antiviral defense: Function and regulation by innate cytokines. *Annual Review of Immunology*, **17**, 189-220.
- Bischoff, S. C. & Kramer, S. (2007). Human mast cells, bacteria, and intestinal immunity. *Immunological Reviews*, **217**, 329-337.
- Bjellqvist, B., Ek, K., Righetti, P. G., Gianazza, E., Gorg, A., Westermeier, R. & Postel, W. (1982). Isoelectric-focusing in immobilized pH gradients - principle, methodology and some applications. *Journal of Biochemical and Biophysical Methods*, **6**, 317-339.
- Bjorck, L. & Kronvall, G. (1984). Purification and some properties of streptococcal Protein-G, a novel IgG-binding reagent. *Journal of Immunology*, **133**, 969-974.
- Blume, U., Ferracin, J., Verschoore, M., Czernielewski, J. M. & Schaefer, H. (1991). Physiology of the vellus hair follicle - hair growth and sebum excretion. *British Journal of Dermatology*, **124**, 21-28.
- Bolduc, C. & Shapiro, J. (2001). The treatment of alopecia areata. *Dermatologic Therapy*, **14**, 306-316.
- Botchkareva, N. V., Ahluwalia, G. & Shander, D. (2006). Apoptosis in the hair follicle. *Journal of Investigative Dermatology*, **126**, 258-264.

- Boyer, R. (2000). *Modern Experimental Biochemistry*. 3rd eds. Addison Wesley Longman, Inc.
- Breathnach, A. S. & Smith, J. (1968). Fine structure of the early hair germ and dermal papilla in the human foetus. *Journal of Anatomy*, **102**, 511-526.
- Brewer, J. W. & Corley, R. B. (1997). Late events in assembly determine the polymeric structure and biological activity of secretory IgM. *Molecular Immunology*, **34**, 323-331.
- Bruet, V., Degorce-Rubiales, F., Abadie, J. & Bourdeau, P. (2008). Severe alopecia areata and onychodystrophy on all four feet of a French trotter mare. *Veterinary Record*, **162**, 758-760.
- Brunda, M. J., Minden, P. & Grey, H. M. (1979). Heterogeneity of binding of human IgA subclasses to protein A. *Journal of Immunology*, **123**, 1457-1461.
- Burnette, W. N. (1981). Western blotting - electrophoretic transfer of proteins from sodium dodecyl sulfate-polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated Protein-A. *Analytical Biochemistry*, **112**, 195-203.
- Bystryn, J. C., Orentreich, N. & Stengel, F. (1979). Direct immunofluorescence studies in alopecia areata and male pattern alopecia. *Journal of Investigative Dermatology*, **73**, 317-320.

- Call, M. E., Pyrdol, J., Wiedmann, M. & Wucherpfennig, K. W. (2002). The organizing principle in the formation of the T cell receptor-CD3 complex. *Cell*, **111**, 967-979.
- Carroll, J. M., McElwee, K. J., King, L. E., Byrne, M. C. & Sundberg, J. P. (2002). Gene array profiling and immunomodulation studies define a cell-mediated immune response underlying the pathogenesis of alopecia areata in a mouse model and humans. *Journal of Investigative Dermatology*, **119**, 392-402.
- Carter, D. M. & Jegasothy, B. V. (1976). Alopecia areata and down syndrome. *Archives of Dermatology*, **112**, 1397-1399.
- Chait, B. T. & Kent, S. B. H. (1992). Weighing naked proteins - practical, high-accuracy mass measurement of peptides and proteins. *Science*, **257**, 1885-1894.
- Chang, S. J. & Lamm, S. H. (2003). Human health effects of sodium azide exposure: A literature review and analysis. *International Journal of Toxicology*, **22**, 175-186.
- Chase, H. B., Rauch, R. & Smith, V. W. (1951). Critical stages of hair development and pigmentation in the mouse. *Physiological Zoology*, **24**, 1-8.
- Chase, H. B. (1954). Growth of the hair. *Physiological Reviews*, **34**, 113-126.
- Chen, C. B. & Wallis, R. (2001). Stoichiometry of complexes between mannose-binding protein and its associated serine proteases - Defining functional

units for complement activation. *Journal of Biological Chemistry*, **276**, 25894-25902.

Chikakane, K. & Takahashi, H. (1995). Measurement of skin pH and its significance in cutaneous diseases. *Clinics in Dermatology*, **13**, 299-306.

Chiodi, F., Siden, A. & Osby, E. (1985). Isoelectric focusing of monoclonal immunoglobulin G, A and M followed by detection with the avidin-biotin system. *Electrophoresis*, **6**, 124-128.

Christoph, T., Müller-Röver, S., Audring, H., Tobin, D. J., Hermes, B., Cotsarelis, G., Ruckert, R. & Paus, R. (2000). The human hair follicle immune system: Cellular composition and immune privilege. *British Journal of Dermatology*, **142**, 862-873.

Chuong, C. M. (1998). Molecular basis of epithelial appendage morphogenesis. Landes Bioscience, Texas.

Colombe, B. W., Price, V. H., Khoury, E. L., Garovoy, M. R. & Lou, C. D. (1995). HLA class-II antigen associations help to define two types of alopecia areata. *Journal of the American Academy of Dermatology*, **33**, 757-764.

Colon, E. A., Popkin, M. K., Callies, A. L., Dessert, N. J. & Hordinsky, M. K. (1991). Lifetime prevalence of psychiatric disorders in patients with alopecia areata. *Comprehensive Psychiatry*, **32**, 245-251.

Cooper, D. & Sun, T. T. (1986). Monoclonal antibody analysis of bovine epithelial keratins - specific pairs as defined by coexpression. *Journal of Biological Chemistry*, **261**, 4646-4654.

- Corbett, B. A., Kantor, A. B., Schulman, H., Walker, W. L., Lit, L., Ashwood, P., Rocke, D. M. & Sharp, F. R. (2007). A proteomic study of serum from children with autism showing differential expression of apolipoproteins and complement proteins. *Molecular Psychiatry*, **12**, 292-306.
- Cotsarelis, G., Sun, T. T. & Lavker, R. M. (1990). Label-Retaining Cells Reside in the Bulge Area of Pilosebaceous Unit - Implications for Follicular Stem-Cells, Hair Cycle, and Skin Carcinogenesis. *Cell*, **61**, 1329-1337.
- Crick, F. H. C. (1953). The packing of alpha-helices: Simple coiled-coils. *Acta Crystallographica*, **6**, 689-697.
- Cui, J., Arita, Y. & Bystryrn, J. C. (1993). Cytolytic Antibodies to Melanocytes in Vitiligo. *Journal of Investigative Dermatology*, **100**, 812-815.
- Cutler, P. (2004a). Bulk purification by fractional precipitation. *In Protein Purification Protocols - Methods in Molecular Biology. Vol. 244*. Human Press Inc, Totowa NJ. 117-124.
- Cutler, P. (2004b). Ion-exchange chromatography. *In Protein Purification Protocols - Methods in Molecular Biology. Vol. 244*. Human Press Inc, Totowa NJ. 125-131.
- Cutler, P. & Doonan, S. (2004a). Affinity chromatography. *In Protein Purification Protocols - Methods in Molecular Biology. Vol. 244*. Human Press Inc, Totowa NJ. 139-149.

- Cutler, P. & Doonan, S. (2004b). General strategies. *In Protein Purification Protocols - Methods in Molecular Biology. Vol. 244.* Human Press Inc, Totowa NJ. 1-13.
- Cytokeratin 16 (C-12): sc-49176 Specifications. Santa Cruz Biotechnology Inc., California.
- Davidsson, P. & Nilsson, C. L. (1999). Peptide mapping of proteins in cerebrospinal fluid utilizing a rapid preparative two-dimensional electrophoretic procedure and matrix-assisted laser desorption/ionization mass spectrometry. *Biochimica et Biophysica Acta-General Subjects*, **1473**, 391-399.
- Davies, A., Simmons, D. L., Hale, G., Harrison, R. A., Tighe, H., Lachmann, P. J. & Waldmann, H. (1989). CD59, an Ly-6-like protein expressed in human lymphoid cells, regulates the action of the complement membrane attack complex on homologous cells. *Journal of Experimental Medicine*, **170**, 637-654.
- Davies, D. R. (1993). Antibody Structure. *Accounts of Chemical Research*, **26**, 421-427.
- Davis, M. M. & Bjorkman, P. J. (1988). T-cell antigen receptor genes and T-cell recognition. *Nature*, **334**, 395-402.
- Degn, S. E., Thiel, S. & Jensenius, J. C. (2007). New perspectives on mannan-binding lectin-mediated complement activation. *Immunobiology*, **212**, 301-311.

- de Hoog, C. L. & Mann, M. (2004). Proteomics. *Annual Review of Genomics & Human Genetics*, **5**, 267-293.
- Ding, X., Aoki, V., Mascaro, J. M., LopezSwiderski, A., Diaz, L. A. & Fairley, J. A. (1997). Mucosal and mucocutaneous (generalized) pemphigus vulgaris show distinct autoantibody profiles. *Journal of Investigative Dermatology*, **109**, 592-596.
- Discipio, R. G. & Gagnon, J. (1982). Characterization of human complement components C6 and C7. *Molecular Immunology*, **19**, 1425-1431.
- Dong, C. (2008). T-H 17 cells in development: an updated view of their molecular identity and genetic programming. *Nature Reviews Immunology*, **8**, 337-348.
- Dovezenski, N., Billetta, R. & Gigli, I. (1992). Expression and Localization of Proteins of the Complement-System in Human Skin. *Journal of Clinical Investigation*, **90**, 2000-2012.
- Dry, F. W. (1926). The coat of the mouse (*Mus Musculus*). *Journal of Genetics*, **165**, 281-340.
- Eckert, J., Church, R. E. & Ebling, F. J. (1968). The pathogenesis of alopecia areata. *British Journal of Dermatology*, **80**, 203-210.
- Edelman, G. M. (1973). Antibody Structure and Molecular Immunology. *Science*, **180**, 830-840.

- Eichner, R., Bonitz, P. & Sun, T. T. (1984). Classification of epidermal keratins according to their immunoreactivity, isoelectric point, and mode of expression. *Journal of Cell Biology*, **98**, 1388-1396.
- Elssner, T., Niemeyer, D., Deininger, S.-O., Ketterlinus, R., Suckau, D., Hahner, S., Jainhuknan, J. & Kostrzewa, M. (2004). Identification of serum peptides by magnetic bead based fractionation combined with LC-MALDI-TOF/TOF. In *HUPO 3 Annual World Congress Beijing*. Bruker Daltonik GmbH, Leipzig, Germany.
- Fairweather, D., Frisancho-Kiss, S. & Rose, N. R. (2008). Sex differences in autoimmune disease from a pathological perspective. *American Journal of Pathology*, **173**, 600-609.
- Fearon, D. T., Austen, K. F. & Ruddy, S. (1974). Properdin Factor D: characterization of its active site and isolation of precursor form. *Journal of Experimental Medicine*, **139**, 355-366.
- Fearon, D. T. & Austen, K. F. (1975). Properdin: Binding to C3b and stabilization of C3b-dependent C3 convertase. *Journal of Experimental Medicine*, **142**, 856-863.
- Field, P. R., Shanker, S. & Murphy, A. M. (1980). The use of Protein A-sepharose affinity chromatography for separation and detection of specific IgM antibody in acquired rubella infection: A comparison with absorption by staphylococci containing Protein-A and density gradient ultracentrifugation. *Journal of Immunological Methods*, **32**, 59-70.
- Fishelson, Z., Attali, G. & Mevorach, D. (2001). Complement and apoptosis. *Molecular Immunology*, **38**, 207-219.

- Fitzpatrick, T. B., Brunet, P. & Kukita, A. (1958). The nature of hair pigment. *In The biology of hair growth*. Montagna, W. & Ellis, R. A. Academic Press Inc., New York. 255-303.
- Fontenot, J. D. & Rudensky, A. Y. (2005). A well adapted regulatory contrivance: regulatory T cell development and the forkhead family transcription factor Foxp3. *Nature Immunology*, **6**, 331-337.
- Forsgren, A. & Sjoquist, J. (1966). "Protein A" from *S. aureus*. I. Pseudo-immune reaction with human gamma-globulin. *Journal of Immunology*, **97**, 822-827.
- Frangion, B., Milstein, C. & Pink, J. R. L. (1969). Structural studies of Immunoglobulin G. *Nature*, **221**, 145-148.
- Franke, W. W., Schiller, D. L., Hatzfeld, M. & Winter, S. (1983). Protein complexes of intermediate-sized filaments - melting of cytoke-
ratin complexes in urea reveals different polypeptide separation characteristics. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences*, **80**, 7113-7117.
- Freyschmidt-Paul, P., Seiter, S., Zoller, M., Konig, A., Ziegler, A., Sundberg, J. P., Happle, R. & Hoffmann, R. (2000). Treatment with an anti-CD44v10-specific antibody inhibits the onset of alopecia areata in C3H/HeJ mice. *Journal of Investigative Dermatology*, **115**, 653-657.
- Freyschmidt-Paul, P., McElwee, K., Botchkarev, V., Sundberg, J., Happle, R. & Hoffmann, R. (2001). The role of Fas and Fas ligand in alopecia areata of C3H/HeJ mice. *Journal of Investigative Dermatology*, **117**, 431-431.

- Fuchs, E. (1995). Keratins and the skin. *Annual Review of Cell & Developmental Biology*, **11**, 123-153.
- Gammon, W. R., Lewis, D. M., Carlo, J. R., Sams, W. M., Jr. & Wheeler, C. E., Jr. (1980). Pemphigoid antibody mediated attachment of peripheral blood leukocytes at the dermal-epidermal junction of human skin. *Journal of Investigative Dermatology*, **75**, 334-339.
- Garcia-Hernandez, M. J., Torres, M. J., Palomares, J. C., Rodriguez-Pichardo, A., Aznar, J. & Camacho, F. (1998). No evidence of cytomegalovirus DNA in alopecia areata. *Journal of Investigative Dermatology*, **110**, 185-185.
- Garn, S. M. (1951). Types and Distribution of the Hair in Man. *Annals of the New York Academy of Sciences*, **53**, 498-507.
- Gasque, P. (2004). Complement: a unique innate immune sensor for danger signals. *Molecular Immunology*, **41**, 1089-1098.
- Gigli, I. & Tausk, F. A. (1988). C1, C4, and C2 components of the classical pathway of complement and regulatory proteins. *Methods in Enzymology*, **162**, 626-639.
- Gilhar, A. & Krueger, G. G. (1987). Hair growth in scalp grafts from patients with alopecia areata and alopecia universalis grafted onto nude mice. *Archives of Dermatology*, **123**, 44-50.
- Gilhar, A., Ullmann, Y., Berkutzki, T., Assy, B. & Kalish, R. S. (1998). Autoimmune hair loss (alopecia areata) transferred by T lymphocytes to

human scalp explants on SCID mice. *Journal of Clinical Investigation*, **101**, 62-67.

Gilhar, A., Shalaginov, R., Assy, B., Serafimovich, S. & Kalish, R. S. (1999). Alopecia areata is a T-lymphocyte mediated autoimmune disease: Lesional human T-lymphocytes transfer alopecia areata to human skin grafts on SCID mice. *Journal of Investigative Dermatology Symposium Proceedings*, **4**, 207-210.

Gilhar, A., Landau, M., Assy, B., Shalaginov, R., Serafimovich, S. & Kalish, R. S. (2001). Melanocyte-associated T cell epitopes can function as autoantigens for transfer of alopecia areata to human scalp explants on Prkdc^{scid} mice. *Journal of Investigative Dermatology*, **117**, 1357-1362.

Gilhar, A., Landau, M., Assy, B., Shalaginov, R., Serafimovich, S. & Kalish, R. S. (2002). Mediation of alopecia areata by cooperation between CD4⁺ and CD8⁺ T lymphocytes - transfer to human scalp explants on Prkdc^{scid} mice. *Archives of Dermatology*, **138**, 916-922.

Gilhar, A. & Kalish, R. S. (2006). Alopecia areata: A tissue specific autoimmune disease of the hair follicle. *Autoimmunity Reviews*, **5**, 64-69.

Gilman, S. (2002). Joint position sense and vibration sense: Anatomical organisation and assessment. *Journal of Neurology Neurosurgery and Psychiatry*, **73**, 473-477.

Gipson, I. K. (1977). Cytoplasmic filaments - their role in motility and cell shape. *Investigative Ophthalmology & Visual Science*, **16**, 1081-1084.

- Goodnow, C. C., Sprent, J., Fazekas de St Groth, B. & Vinuesa, C. G. (2005). Cellular and genetic mechanisms of self tolerance and autoimmunity. *Nature*, **435**, 590-597.
- Goodnow, C. C. (2007). Multistep pathogenesis of autoimmune disease. *Cell*, **130**, 25-35.
- Gorovsky, M. A., Carlson, K. & Rosenbaum, J. L. (1970). Simple method for quantitative densitometry of polyacrylamide gels using fast green. *Analytical Biochemistry*, **35**, 359-370.
- Gorski, J. P., Hugli, T. E. & Müller-Eberhard, H. J. (1979). C4a - The third anaphylatoxin of the human complement system. *Proceedings of the National Academy of Sciences of the United States of America*, **76**, 5299-5302.
- Gu, L. H. & Coulombe, P. A. (2007). Keratin expression provides novel insight into the morphogenesis and function of the companion layer in hair follicles. *Journal of Investigative Dermatology*, **127**, 1061-1073.
- Gupta, A. K., Ellis, C. N., Cooper, K. D., Nickoloff, B. J., Ho, V. C., Chan, L. S., Hamilton, T. A., Tellner, D. C., Griffiths, C. E. M. & Voorhees, J. J. (1990). Oral cyclosporine for the treatment of alopecia areata - a clinical and immunohistochemical analysis. *Journal of the American Academy of Dermatology*, **22**, 242-250.
- Gygi, S. P., Corthals, G. L., Zhang, Y., Rochon, Y. & Aebersold, R. (2000). Evaluation of two-dimensional gel electrophoresis-based proteome analysis technology. *Proceedings of the National Academy of Sciences of the United States of America*, **97**, 9390-9395.

- Habeeb, A. F. & Francis, R. D. (1976). Preparation of human immunoglobulin by ammonium sulfate precipitation. *Vox Sanguinis*, **31**, 423-434.
- Hamm, H., Klemmer, S., Kreuzer, I., Steijlen, P. M., Happler, R. & Brocker, E. B. (1988). HLA-DR and HLA-DQ antigen expression of anagen and telogen hair bulbs in long-standing alopecia areata. *Archives of Dermatological Research*, **280**, 179-181.
- Hardy, G., Bevan, S. J., McElroy, B., Palmer, T. E. A., Griffiths, R. D. & Braidwood, C. (1993). Stability of glutamine in parenteral feeding solutions. *Lancet*, **342**, 186-186.
- Harty, J. T., Tvinnereim, A. R. & White, D. W. (2000). CD8(+) T cell effector mechanisms in resistance to infection. *Annual Review of Immunology*, **18**, 275-308.
- Harwood, N. E. & Batista, F. D. (2008). New insights into the early molecular events underlying B cell activation. *Immunity*, **28**, 609-619.
- Hatzfeld, M. & Franke, W. W. (1985). Pair formation and promiscuity of cytokeratins - formation in vitro of heterotypic complexes and intermediate-sized filaments by homologous and heterologous recombinations of purified polypeptides. *Journal of Cell Biology*, **101**, 1826-1841.
- Headington, J. T. (1984). Transverse microscopic anatomy of the human scalp - a basis for a morphometric approach to disorders of the hair follicle. *Archives of Dermatology*, **120**, 449-456.

- Headington, J. T. (1991). The histopathology of alopecia areata. *Journal of Investigative Dermatology*, **96**, S69-S69.
- Hedstrand, H., Perheentupa, J., Ekwall, O., Gustafsson, J., Michaelsson, G., Husebye, E., Rorsman, F. & Kampe, O. (1999). Antibodies against hair follicles are associated with alopecia totalis in autoimmune polyendocrine syndrome type I. *Journal of Investigative Dermatology*, **113**, 1054-1058.
- Herrmann, H. & Aebi, U. (2004). Intermediate filaments: Molecular structure, assembly mechanism, and integration into functionally distinct intracellular scaffolds. *Annual Review of Biochemistry*, **73**, 749-789.
- Holmskov, U., Thiel, S. & Jensenius, J. C. (2003). Collectins and ficolins: Humoral lectins of the innate immune defense. *Annual Review of Immunology*, **21**, 547-578.
- Hordinsky, M. & Ericson, M. (2004). Autoimmunity: Alopecia areata. *Journal of Investigative Dermatology Symposium Proceedings*, **9**, 73-78.
- Inazu, M. & Mishima, Y. (1993). Detection of eumelanogenic and pheomelanogenic melanosomes in the same normal human melanocyte. *Journal of Investigative Dermatology*, **100**, S172-S175.
- Ito, M. (1990). Dynamic Ultrastructural-Changes of the Connective-Tissue Sheath of Human Hair-Follicles During Hair Cycle. *Archives of Dermatological Research*, **282**, 434-441.
- Jackow, C., Puffer, N., Hordinsky, M., Nelson, J., Tarrand, J. & Duvic, M. (1998). Alopecia areata and cytomegalovirus infection in twins: Genes versus

environment? *Journal of the American Academy of Dermatology*, **38**, 418-425.

Jackson, D., Church, R. E. & Ebling, F. J. (1971). Alopecia areata hairs - scanning electron microscope study. *British Journal of Dermatology*, **85**, 242-246.

Jahoda, C. A. B., Horne, K. A. & Oliver, R. F. (1984). Induction of hair growth by implantation of cultured dermal papilla cells. *Nature*, **311**, 560-562.

Jaks, V. (in press). Lgr5 marks dynamic hair follicle stem cells. *Nature Genetics*.

Jarrett, A. (1977). The Hair Follicle. eds. Vol. 4. Academic Press Inc., London.

Jimbow, K., Park, J. S., Kato, F., Hirosaki, K., Toyofuku, K., Hua, C. & Yamashita, T. (2000). Assembly, target-signaling and intracellular transport of tyrosinase gene family proteins in the initial stage of melanosome biogenesis. *Pigment Cell Research*, **13**, 222-229.

Johnson, R. S., Martin, S. A., Biemann, K., Stults, J. T. & Watson, J. T. (1987). Novel fragmentation process of peptides by collision-induced decomposition in a tandem mass spectrometer: differentiation of leucine and isoleucine. *Analytical Chemistry*, **59**, 2621-2625.

Kanno, T., Kawada, A., Yamanouchi, J., Yosida-Noro, C., Yoshiki, A., Shiraiwa, M., Kusakabe, M., Manabe, M., Tezuka, T. & Takahara, H. (2000). Human peptidylarginine deiminase type III: Molecular cloning and nucleotide sequence of the cDNA, properties of the recombinant enzyme,

and immunohistochemical localization in human skin. *Journal of Investigative Dermatology*, **115**, 813-823.

Kaplan, A. P. (2005). Does C-2 kinin exist? *Journal of Allergy & Clinical Immunology*, **115**, 876-876.

Kapsenberg, M. L. (2003). Dendritic-cell control of pathogen-driven T-cell polarization. *Nature Reviews Immunology*, **3**, 984-993.

Kavak, A., Baykal, C., Ozarmagan, G. & Akar, U. (2000). HLA in alopecia areata. *International Journal of Dermatology*, **39**, 589-592.

Kawasaki, N., Kawasaki, T. & Yamashina, I. (1989). A serum lectin (mannan-binding protein) has complement-dependent bactericidal activity. *Journal of Biochemistry*, **106**, 483-489.

Kemp, E. H., McDonagh, A. J. G., Wengraf, D. A., Messenger, A. G., Gawkrödger, D. J., Cork, M. J. & Tazi-Ahnini, R. (2006). The non-synonymous C1858T substitution in the PTPN22 gene is associated with susceptibility to the severe forms of alopecia areata. *Human Immunology*, **67**, 535-539.

Khoury, E. L., Price, V. H. & Greenspan, J. S. (1988). HLA-DR expression by hair follicle keratinocytes in alopecia areata - evidence that it is secondary to the lymphoid infiltration. *Journal of Investigative Dermatology*, **90**, 193-200.

- Ki67 Antigen Ready to Use Liquid Mouse Monoclonal Antibody Instructions for Use. (2004). Novocastra Laboratories Ltd., Newcastle Upon Tyne. RTU-Ki67-MM1/CE/UK, Form 772 rev- 5/08/03.
- Kimura, Y., Miwa, T., Zhou, L. & Song, W.-C. (2008). Activator-specific requirement of properdin in the initiation and amplification of the alternative pathway complement. *Blood*, **111**, 732-740.
- King Jr., L. E., McElwee, K. J. & Sundberg, J. P. (2008). Alopecia areata. *In Current Directions in Autoimmunity. Vol. 10*. Edited by Nickoloff, B. J. & Nestle, F. O. 280-312.
- Kinoshita, T., Takata, Y., Kozono, H., Takeda, J., Hong, K. & Inoue, K. (1988). C5 convertase of the alternative complement pathway: covalent linkage between two C3b molecules within the trimolecular complex enzyme. *Journal of Immunology*, **141**, 3895-3901.
- Kitsioulis, E., Lekka, M. E., Nakos, G., Cassagne, C. & Maneta-Peyret, L. (2002). Lipids are co-eluted with immunoglobulins G during purification by recombinant streptococcal Protein-G affinity chromatography. *Journal of Immunological Methods*, **271**, 107-111.
- Klickstein, L. B., Bartow, T. J., Miletic, V., Rabson, L. D., Smith, J. A. & Fearon, D. T. (1988). Identification of distinct C3b and C4b recognition sites in the human C3b/C4b receptor (CR1, CD35) by deletion mutagenesis. *Journal of Experimental Medicine*, **168**, 1699-1717.
- Kligman, A. M. (1959). The human hair cycle. *Journal of Investigative Dermatology*, **33**, 307-316.

- Kolb, W. P., Haxby, J. A., Arroyave, C. M. & Müller-Eberhard, H. J. (1972). Molecular analysis of membrane attack mechanism of complement. *Journal of Experimental Medicine*, **135**, 549-566.
- Kraal, G., Weissman, I. L. & Butcher, E. C. (1982). Germinal Center B-Cells - Antigen-Specificity and Changes in Heavy-Chain Class Expression. *Nature*, **298**, 377-379.
- Kronvall, G. & Williams, R. C., Jr. (1969). Differences in anti-Protein-A activity among IgG subgroups. *Journal of Immunology*, **103**, 828-833.
- Lachgar, S., Charveron, M., Gall, Y. & Bonafe, J. L. (1998). Minoxidil upregulates the expression of vascular endothelial growth factor in human hair dermal papilla cells. *British Journal of Dermatology*, **138**, 407-411.
- Lachmann, P. J. & Halbwachs, L. (1975). Influence of C3b inactivator (KAF) concentration on ability of serum to support complement activation. *Clinical & Experimental Immunology*, **21**, 109-114.
- Langbein, L., Rogers, M. A., Winter, H., Praetzel, S., Beckhaus, U., Rackwitz, H. R. & Schweizer, J. (1999). The catalog of human hair keratins I. Expression of the nine type I members in the hair follicle. *Journal of Biological Chemistry*, **274**, 19874-19884.
- Langbein, L., Rogers, M. A., Winter, H., Praetzel, S. & Schweizer, J. (2001). The catalog of human hair keratins. II. Expression of the six type II members in the hair follicle and the combined catalog of human type I and II keratins. *Journal of Biological Chemistry*, **276**, 35123-35132.

- Langbein, L., Rogers, M. A., Praetzel, S., Winter, H. & Schweizer, J. (2003). K6irs1, K6irs2, K6irs3, and K6irs4 represent the inner-root-sheath-specific type II epithelial keratins of the human hair follicle. *Journal of Investigative Dermatology*, **120**, 512-522.
- Langbein, L., Rogers, M. A., Praetzel-Wunder, S., Helmke, B., Schirmacher, P. & Schweizer, J. (2006). K25 (K25irs1), K26 (K25irs2), K27 (K25irs3), and K28 (K25irs4) represent the type I inner root sheath keratins of the human hair follicle. *Journal of Investigative Dermatology*, **126**, 2377-2386.
- Lanzavecchia, A. (1985). Antigen specific interaction between T cells and B cells. *Nature*, **314**, 537-539.
- Lee, S. C., Kim, I. G., Marekov, L. N., Okeefe, E. J., Parry, D. A. D. & Steinert, P. M. (1993). The Structure of Human Trichohyalin - Potential Multiple Roles as a Functional Ef-Hand-Like Calcium-Binding Protein, a Cornified Cell-Envelope Precursor, and an Intermediate Filament-Associated (Cross-Linking) Protein. *Journal of Biological Chemistry*, **268**, 12164-12176.
- Lefranc, D., Almeras, L., Dubucquoi, S., de Seze, J., Vermersch, P. & Prin, L. (2004). Distortion of the self-reactive IgG antibody repertoire in multiple sclerosis as a new diagnostic tool. *Journal of Immunology*, **172**, 669-678.
- Lejeune, J. & Turpin, R. (1961). Chromosomal aberrations in Man. *American Journal of Human Genetics*, **13**, 175-184.
- Leon, B. & Ardavin, C. (2008). Monocyte-derived dendritic cells in innate and adaptive immunity. *Immunology & Cell Biology*, **86**, 320-324.

- Li, N., Aoki, V., Hans, G., Rivitti, E. A. & Diaz, L. A. (2003). The role of intramolecular epitope spreading in the pathogenesis of endemic pemphigus foliaceus (Fogo Selvagem). *Journal of Experimental Medicine*, **197**, 1501-1510.
- Lindner, G., Botchkarev, V. A., Botchkareva, N. V., Ling, G., van der Veen, C. & Paus, R. (1997). Analysis of apoptosis during hair follicle regression (catagen). *American Journal of Pathology*, **151**, 1601-1617.
- Loser, K. & Beissert, S. (2007). Dendritic cells and T cells in the regulation of cutaneous immunity. *Advances in Dermatology*, **23**, 307-333.
- Lowry, O. H., Rosenbrough, N. J., Farr, A. L. & Randall, R. J. (1951). Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry*, **193**, 265-275.
- Lublin, D. M. & Atkinson, J. P. (1989). Decay-accelerating factor - Biochemistry, molecular biology, and function. *Annual Review of Immunology*, **7**, 35-58.
- Lueking, A., Huber, O., Wirths, C., Schulte, K., Stieler, K. M., Blume-Peytavi, U., Kowald, A., Hensel-Wiegel, K., Tauber, R., Lehrach, H., Meyer, H. E. & Cahill, D. J. (2005). Profiling of alopecia areata autoantigens based on protein microarray technology. *Molecular & Cellular Proteomics*, **4**, 1382-1390.
- Lyle, S., Christofidou-Solomidou, M., Liu, Y. P., Elder, D. E., Albelda, S. & Cotsarelis, G. (1998). The C8/144B monoclonal antibody recognizes cytokeratin 15 and defines the location of human hair follicle stem cells. *Journal of Cell Science*, **111**, 3179-3188.

- Lynch, M. H., Oguin, W. M., Hardy, C., Mak, L. & Sun, T. T. (1986). Acidic and basic hair nail (hard) keratins - their colocalization in upper cortical and cuticle cells of the human hair follicle and their relationship to soft keratins. *Journal of Cell Biology*, **103**, 2593-2606.
- Madani, S. & Shapiro, J. (2000). Alopecia areata update. *Journal of the American Academy of Dermatology*, **42**, 549-566.
- Magerl, M., Kausler, S., Paus, R. & Tobin, D. J. (2002). Simple and rapid method to isolate and culture follicular papillae from human scalp hair follicles. *Experimental Dermatology*, **11**, 381-385.
- Magin, T. M., Vijayaraj, P. & Leube, R. E. (2007). Structural and regulatory functions of keratins. *Experimental Cell Research*, **313**, 2021-2032.
- Mahoney, M. G., Wang, Z., Rothenberger, K., Koch, P. J., Amagai, M. & Stanley, J. R. (1999). Explanations for the clinical and microscopic localization of lesions in pemphigus foliaceus and vulgaris. *Journal of Clinical Investigation*, **103**, 461-468.
- Manabe, M. & O'Guin, W. M. (1994). Existence of trichohyalin-keratohyalin hybrid granules - colocalization of two major intermediate filament-associated proteins in non-follicular epithelia. *Differentiation*, **58**, 65-75.
- Mancini, G., Carbonara, A. O. & Heremans, J. F. (1965). Immunochemical quantitation of antigens by single radial immunodiffusion. *Immunochemistry*, **2**, 235-254.

- Mangelsdorf, S., Otberg, N., Maibach, H. I., Sinkgraven, R. & Sterry, W. (2006). Ethnic variation in vellus hair follicle size and distribution. *Skin Pharmacology & Physiology*, **19**, 159-167.
- Mann, M., Hendrickson, R. C. & Pandey, A. (2001). Analysis of proteins and proteomes by mass spectrometry. *Annual Review of Biochemistry*, **70**, 437-473.
- Martinez-Mir, A., Zlotogorski, A., Gordon, D., Petukhova, L., Mo, J., Gilliam, T. C., Londono, D., Haynes, C., Ott, J., Hordinsky, M., Nanova, K., Norris, D., Price, V., Duvic, M. & Christiano, A. M. (2007). Genomewide scan for linkage reveals evidence of several susceptibility loci for alopecia areata. *American Journal of Human Genetics*, **80**, 316-328.
- Matsudaira, P. (1987). Sequence from picomole quantities of proteins electroblotted onto polyvinylidene difluoride membranes. *Journal of Biological Chemistry*, **262**, 10035-10038.
- Matsushita, M., Endo, Y. & Fujita, T. (2000). Cutting edge: Complement-activating complex of ficolin and mannose-binding lectin-associated serine protease. *Journal of Immunology*, **164**, 2281-2284.
- McDonagh, A. J. G. & Tazi-Ahnini, R. (2002). Epidemiology and genetics of alopecia areata. *Clinical & Experimental Dermatology*, **27**, 405-409.
- McElwee, K. J., Spiers, E. M. & Oliver, R. F. (1996). In vivo depletion of CD8⁺ T cells restores hair growth in the DEBR model for alopecia areata. *British Journal of Dermatology*, **135**, 211-217.

- McElwee, K. J., Boggess, D., Burgett, B., Bates, R., Bedigan, H. G., Sundberg, J. P. & King, L. E. (1998). Murine cytomegalovirus is not associated with alopecia areata in C3H/HeJ mice. *Journal of Investigative Dermatology*, **110**, 986-987.
- McElwee, K. J., Spiers, E. M. & Oliver, R. F. (1999). Partial restoration of hair growth in the DEBR model for alopecia areata after in vivo depletion of CD4⁺ T cells. *British Journal of Dermatology*, **140**, 432-437.
- McElwee, K. J., Hoffmann, R., Freyschmidt-Paul, P., Wenzel, E., Kissling, S., Sundberg, J. P. & Zoller, M. (2002). Resistance to alopecia areata in C3H/HeJ mice is associated with increased expression of regulatory cytokines and a failure to recruit CD4⁺ and CD8⁺ cells. *Journal of Investigative Dermatology*, **119**, 1426-1433.
- McElwee, K. J., Freyschmidt-Paul, P., Hoffmann, R., Kissling, S., Hummel, S., Vitacolonna, M. & Zoller, M. (2005). Transfer of CD8⁺ cells induces localized hair loss whereas CD4⁺/CD25⁻ cells promote systemic alopecia areata and CD4⁺/CD25⁺ cells blockade disease onset in the C3H/HeJ mouse model. *Journal of Investigative Dermatology*, **124**, 947-957.
- McPhee, C., Kennedy, V., Silva, K. A., King, B., King, L. E., Roopenian, D. & Sundberg, J. P. (2008). Testing human alopecia areata candidate genes in the C3H/HeJ mouse model system. *Journal of Investigative Dermatology*, **128**, S117-S117.
- Medzhitov, R. & Janeway, C. A. (1997). Innate immunity: The virtues of a nonclonal system of recognition. *Cell*, **91**, 295-298.

- Messenger, A. G., Slater, D. N. & Bleehen, S. S. (1986). Alopecia areata - alterations in the hair growth cycle and correlation with the follicular pathology. *British Journal of Dermatology*, **114**, 337-347.
- Metschnikoff, E. (1884). Ueber eine Sprosspilzkrankheit der Daphnien. Beitrag zur Lehre über den Kampf der Phagocyten gegen Krankheitserreger, Archiv f. pathologische Anatomie und Physiologie und f. klinische Medizin, 96: 177-195. *In Milestones in Microbiology: 1556 to 1940*. Translated and edited by Brock, T. D. ASM Press, 1998, P.132.
- Metz, M., Siebenhaar, F. & Maurer, M. (2008). Mast cell functions in the innate skin immune system. *Immunobiology*, **213**, 251-260.
- Meyer, T. S. & Lamberts, B. L. (1965). Use of coomassie brilliant blue R250 for the electrophoresis of microgram quantities of parotid saliva proteins on acrylamide-gel strips. *Biochimica et Biophysica Acta*, **107**, 144-145.
- Michie, H. J., Jahoda, C. A. B., Oliver, R. F. & Johnson, B. E. (1991). The DEBR rat: an animal model of human alopecia areata. *British Journal of Dermatology*, **125**, 94-100.
- Miletic, V. D. & Frank, M. M. (1995). Complement Immunoglobulin Interactions. *Current Opinion in Immunology*, **7**, 41-47.
- Moll, R., Franke, W. W., Schiller, D. L., Geiger, B. & Krepler, R. (1982). The catalog of human cytokeratins - patterns of expression in normal epithelia, tumors and cultured cells. *Cell*, **31**, 11-24.

- Moll, R., Divo, M. & Langbein, L. (2008). The human keratins: biology and pathology. *Histochemistry & Cell Biology*, **129**, 705-733.
- Montage® Antibody Purification Kit and Spin Columns with PROSEP®-G Media User Guide (2004). Millipore Corporation, Massachusetts, P36491, Rev. C, 09/04.
- Montagna, W. & van Scott, E. J. (1958). The anatomy of the hair follicle. *In The biology of hair growth*. Montagna, W. & Ellis, R. A. Academic Press Inc., New York. 39-63.
- Morgan, B. P. (1989). Complement membrane attack on nucleated cells - Resistance, recovery and non-lethal effects. *Biochemical Journal*, **264**, 1-14.
- Morling, N., Frenzt, G., Fugger, L., Georgsen, J., Jakobsen, B., Odum, N. & Svejgaard, A. (1991). DNA polymorphism of HLA class II genes in alopecia areata. *Disease Markers*, **9**, 35-42.
- Moser, B., Loetscher, M., Piali, L. & Loetscher, P. (1998). Lymphocyte responses to chemokines. *International Reviews of Immunology*, **16**, 323-344.
- Mouse Anti-Cytokeratin AE1/AE3 Monoclonal Antibody Specifications (2004). Chemicon International, Millipore Corporation, Massachusetts, 1/19/2004/MAB3412/KDL/030805/ef.
- Müller-Eberhard, H. J., Polley, M. J. & Calcott, M. A. (1967). Formation and functional significance of a molecular complex derived from second and

fourth component of human complement. *Journal of Experimental Medicine*, **125**, 359-380.

Müller-Eberhard, H. J. & Gotze, O. (1972). C3 proactivator convertase and its mode of action. *Journal of Experimental Medicine*, **135**, 1003-1008.

Müller-Eberhard, H. J. (1986). The membrane attack complex of complement. *Annual Review of Immunology*, **4**, 503-528.

Müller-Röver, S., Handjiski, B., van der Veen, C., Eichmüller, S., Foitzik, K., McKay, I. A., Stenn, K. S. & Paus, R. (2001). A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *Journal of Investigative Dermatology*, **117**, 3-15.

Murphy, K., Travers, P. & Walport, M. (2008). *Janeway's Immunobiology*. 7th eds. Garland Science, New York.

Myhre, E. B. & Kronvall, G. (1980). Demonstration of specific binding sites for human serum albumin in group C and G streptococci. *Infection & Immunity*, **27**, 6-14.

Nagamine, K., Peterson, P., Scott, H. S., Kudoh, J., Minoshima, S., Heino, M., Krohn, K. J. E., Lalioti, M. D., Mullis, P. E., Antonarakis, S. E., Kawasaki, K., Asakawa, S., Ito, F. & Shimizu, N. (1997). Positional cloning of the APECED gene. *Nature Genetics*, **17**, 393-398.

Nagasawa, S. & Stroud, R. M. (1977). Cleavage of C2 by C1s into antigenically distinct fragments C2a and C2b: Demonstration of binding of C2b to C4b.

Proceedings of the National Academy of Sciences of the United States of America, **74**, 2998-3001.

Nestle, F. O., Zheng, X. G., Thompson, C. B., Turka, L. A. & Nickoloff, B. J. (1993). Characterization of dermal dendritic cells obtained from normal human skin reveals phenotypic and functionally distinctive subsets. *Journal of Immunology*, **151**, 6535-6545.

Nicholson-weller, A., Burge, J., Fearon, D. T., Weller, P. F. & Austen, K. F. (1982). Isolation of a human erythrocyte membrane glycoprotein with decay-accelerating activity for C3 convertases of the complement system. *Journal of Immunology*, **129**, 184-189.

Nishimura, E. K., Jordan, S. A., Oshima, H., Yoshida, H., Osawa, M., Moriyama, M., Jackson, I. J., Barrandon, Y., Miyachi, Y. & Nishikawa, S. (2002). Dominant role of the niche in melanocyte stem-cell fate determination. *Nature*, **416**, 854-860.

Norris, D. A., Kissinger, R. M., Naughton, G. M. & Bystry, J. C. (1988). Evidence for Immunological Mechanisms in Human Vitiligo - Patients Sera Induce Damage to Human Melanocytes In vitro by Complement-Mediated Damage and Antibody-Dependent Cellular Cyto-Toxicity. *Journal of Investigative Dermatology*, **90**, 783-789.

Norris, D. A. (2003). How close are we to solving the puzzle? Review of the alopecia areata research workshop David Norris. *Journal of Investigative Dermatology Symposium Proceedings*, **8**, 222-225.

O'Farrell, P. H. (1975). High resolution two-dimensional electrophoresis of proteins. *Journal of Biological Chemistry*, **250**, 4007-4021.

- O'Garra, A., Barrat, F. J., Castro, A. G., Vicari, A. & Hawrylowicz, C. (2008). Strategies for use of IL-10 or its antagonists in human disease. *Immunological Reviews*, **223**, 114-131.
- O'Guin, W. M., Sun, T. T. & Manabe, M. (1992). Interaction of trichohyalin with intermediate filaments: Three immunologically defined stages of trichohyalin maturation. *Journal of Investigative Dermatology*, **98**, 24-32.
- O'Shea, M. L. (2003). IgG-isotype subclass-usage by the autoantibody response in the human autoimmune hair loss disorder alopecia areata. MSc Dissertation, University of Bradford, UK.
- Okumura, S., Kashiwakura, J., Tomita, H., Matsumoto, K., Nakajima, T., Saito, H. & Okayama, Y. (2003). Identification of specific gene expression profiles in human mast cells mediated by Toll-like receptor 4 and Fc epsilon RI. *Blood*, **102**, 2547-2554.
- Olsen, E., Hordinsky, M., McDonald-Hull, S., Price, V., Roberts, J., Shapiro, J. & Stenn, K. (1999). Alopecia areata investigational assessment guidelines. *Journal of the American Academy of Dermatology*, **40**, 242-246.
- Olsen, E. A. (2003). Disorders of Hair Growth: Diagnosis and Treatment. 2nd eds. McGraw-Hill Companies, Inc.
- Olsen, E. A., Hordinsky, M. K., Price, V. H., Roberts, J. L., Shapiro, J., Canfield, D., Duvic, M., King, L. E., McMichael, A. J., Randall, V. A., Turner, M. L., Sperling, L., Whiting, D. A. & Norris, D. (2004). Alopecia areata investigational assessment guidelines - part II. *Journal of the American Academy of Dermatology*, **51**, 440-447.

- Otberg, N., Richter, H., Schaefer, H., Blume-Peytavi, U., Sterry, W. & Lademann, J. (2004). Variations of hair follicle size and distribution in different body sites. *Journal of Investigative Dermatology*, **122**, 14-19.
- Page, M. & Thorpe, R. (1996). Purification of IgG using DEAE-sepharose chromatography. *In The Protein Protocols Handbook*. Walker, J. M. Humana Press Inc., Totowa, NJ. 725-726.
- Pangburn, M. K., Schreiber, R. D. & Müller-Eberhard, H. J. (1977). Human complement C3b inactivator: isolation, characterization, and demonstration of an absolute requirement for serum protein Beta-1h for cleavage of C3b and C4b in solution. *Journal of Experimental Medicine*, **146**, 257-270.
- Pangburn, M. K., Schreiber, R. D. & Müller-Eberhard, H. J. (1981). Formation of the initial C3 convertase of the alternative complement pathway - acquisition of C3b-like activities by spontaneous hydrolysis of the putative thioester in native C3. *Journal of Experimental Medicine*, **154**, 856-867.
- Pangburn, M. K. & Rawal, N. (2002). Structure and function of complement C5 convertase enzymes. *Biochemical Society Transactions*, **30**, 1006-1010.
- Paradis, M., Fecteau, G. & Scott, D. W. (1988). Alopecia areata (Pelade) in a cow. *Canadian Veterinary Journal-Revue Veterinaire Canadienne*, **29**, 727-729.
- Parker, D. C. (1993). T cell - Dependent B cell activation. *Annual Review of Immunology*, **11**, 331-360.

- Parker, K. C., Garrels, J. I., Hines, W., Butler, E. M., McKee, A. H. Z., Patterson, D. & Martin, S. (1998). Identification of yeast proteins from two-dimensional gels: Working out spot cross-contamination. *Electrophoresis*, **19**, 1920-1932.
- Pasch, M. C., van den Bosch, N. H. A., Daha, M. R., Bos, J. D. & Asghar, S. S. (2000). Synthesis of complement components C3 and factor B in human keratinocytes is differentially regulated by cytokines. *Journal of Investigative Dermatology*, **114**, 78-82.
- Pauling, L., Corey, R. B. & Branson, H. R. (1951). The structure of proteins: Two hydrogen-bonded helical configurations of the polypeptide chain. *Proceedings of the National Academy of Sciences of the United States of America*, **37**, 205-211.
- Paus, R., Handjiski, B., Czarnetzki, B. M. & Eichmuller, S. (1994). A murine model for inducing and manipulating hair follicle regression (catagen) - effects of dexamethasone and cyclosporine-A. *Journal of Investigative Dermatology*, **103**, 143-147.
- Paus, R., Muller-Rover, S., van der Veen, C., Maurer, M., Eichmuller, S., Ling, G., Hofmann, U., Foitzik, K., Mecklenburg, L. & Handjiski, B. (1999). A comprehensive guide for the recognition and classification of distinct stages of hair follicle morphogenesis. *Journal of Investigative Dermatology*, **113**, 523-532.
- Paus, R., Ito, N., Takigawa, M. & Ito, T. (2003). The hair follicle and immune privilege. *Journal of Investigative Dermatology Symposium Proceedings*, **8**, 188-194.

- Paus, R., Ito, N. & Ito, T. (2004). The theory of immune privilege of the hair follicle. *In Animal Models of Human Inflammatory Skin Diseases*. Edited by Chan, L. S. CRC Press. 155-165.
- Paus, R., Nickoloff, B. J. & Ito, T. (2005). A 'hairy' privilege. *Trends in Immunology*, **26**, 32-40.
- Peggs, K. S. & Allison, J. P. (2005). Co-stimulatory pathways in lymphocyte regulation: the immunoglobulin superfamily. *British Journal of Haematology*, **130**, 809-824.
- Peng, J. M. & Gygi, S. P. (2001). Proteomics: The move to mixtures. *Journal of Mass Spectrometry*, **36**, 1083-1091.
- Perkins, D. N., Pappin, D. J. C., Creasy, D. M. & Cottrell, J. S. (1999). Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis*, **20**, 3551-3567.
- Perret, C., Wiesnermenzel, L. & Happle, R. (1984). Immunohistochemical analysis of T cell subsets in the peribulbar and intrabulbar infiltrates of alopecia areata. *Acta Dermato-Venereologica*, **64**, 26-30.
- Person, M. D., Shen, J., Traner, A., Hensley, S. C., Lo, H. H., Abbruzzese, J. L. & Li, D. (2006). Protein fragment domains identified using 2D gel electrophoresis/MALDI-TOF. *Journal of Biomolecular Techniques*, **17**, 145-56.
- Peters, E. M., Liotiri, S., Bodo, E., Ohnemus, U., Biro, T., Arck, P. C. & Paus, R. (2005). Probing the effects of "stress" on the human hair follicle:

Substance P upregulates MHC class I and NGF immunoreactivity ex vivo and stimulates degranulation of connective tissue sheath mast cells. *Journal of Investigative Dermatology*, **125**, A44-A44.

Philpott, M. P., Sanders, D., Westgate, G. E. & Kealey, T. (1994). Human hair growth in vitro: a model for the study of hair follicle biology. *Journal of Dermatological Science*, **7 Suppl**, S55-S72.

Philpott, M. P., Sanders, D. A. & Kealey, T. (1996). Whole hair follicle culture. *Dermatologic Clinics*, **14**, 595-607.

Pober, J. S., Kluger, M. S. & Schechner, J. S. (2001). Human endothelial cell presentation of antigen and the homing of memory/effector T cells to skin. *Cutaneous T Cell Lymphoma: Basic & Clinically Relevant Biology*, **941**, 12-25.

Podack, E. R. & Tschopp, J. (1982). Polymerization of the 9th component of complement (C9): formation of poly(C9) with a tubular ultrastructure resembling the membrane attack complex of complement. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences*, **79**, 574-578.

Price, V. H. (1991). Alopecia areata - clinical aspects. *Journal of Investigative Dermatology*, **96**, S68-S68.

Price, V. H. & Khoury, E. L. (1991). Alopecia Areata. *In Progress in Dermatology*. 25. Moshell, A. N. Dermatology Foundation, Evanston, Illinois. 1-8.

Prota, G. (1980). Recent Advances in the Chemistry of Melanogenesis in Mammals. *Journal of Investigative Dermatology*, **75**, 122-127.

Protein G Immobilized On Agarose Product Information, P-3296 Protein G on Sepharose® 4B fast flow (1998). Sigma-Aldrich Company Ltd., Dorset, P7700, 11/23/98 – LMY

Punnonen, J. & de Vries, J. E. (1994). IL-13 induces proliferation, Ig isotype switching, and Ig synthesis by immature human fetal B cells. *Journal of Immunology*, **152**, 1094-1102.

Ranki, A., Kianto, U., Kanerva, L., Tolvanen, E. & Johansson, E. (1984). Immunohistochemical and electron microscopic characterization of the cellular infiltrate in alopecia (areata, totalis, and universalis). *Journal of Investigative Dermatology*, **83**, 7-11.

Recht, B., Frangione, B., Franklin, E. & van Loghem, E. (1981). Structural studies of a human gamma 3 myeloma protein (Goe) that binds staph Protein-A. *Journal of Immunology*, **127**, 917-923.

Rogers, G. E., Fietz, M. J. & Fratini, A. (1991). Trichohyalin and Matrix Proteins. *Annals of the New York Academy of Sciences*, **642**, 64-81.

Rogers, M. A., Winter, H., Langbein, L., Wolf, C. & Schweizer, J. (2000). Characterization of a 300 kbp region of human DNA containing the type II hair keratin gene domain. *Journal of Investigative Dermatology*, **114**, 464-472.

Rogers, M. A., Winter, H., Langbein, L., Bleiler, R. & Schweizer, J. (2004). The human type I keratin gene family: Characterization of new hair follicle

specific members and evaluation of the chromosome 17q21.2 gene domain. *Differentiation*, **72**, 527-540.

Rokhsar, C. K., Shupack, J. L., Vafai, J. J. & Washenik, K. (1998). Efficacy of topical sensitizers in the treatment of alopecia areata. *Journal of the American Academy of Dermatology*, **39**, 751-761.

Rollins, S. A. & Sims, P. J. (1990). The complement-inhibitory activity of CD59 resides in its capacity to block incorporation of C9 into membrane C5b-9. *Journal of Immunology*, **144**, 3478-3483.

Rook, A. & Dawber, R. (1991). *Diseases of the Hair and Scalp*. 2nd eds. Blackwell Scientific Publications, Oxford.

Rosen, A. & Casciola-Rosen, L. (2001). Clearing the way to mechanisms of autoimmunity. *Nature Medicine*, **7**, 664-665.

Rothnagel, J. A. & Rogers, G. E. (1986). Trichohyalin, an intermediate filament-associated protein of the hair follicle. *Journal of Cell Biology*, **102**, 1419-1429.

Ruiz-Doblado, S., Carrizosa, A. & Garcia-Hernandez, M. J. (2003). Alopecia areata: psychiatric comorbidity and adjustment to illness. *International Journal of Dermatology*, **42**, 434-437.

Safavi, K. H., Muller, S. A., Suman, V. J., Moshell, A. N. & Melton, L. J. (1995). Incidence of alopecia areata in Olmsted County, Minnesota, 1975 through 1989. *Mayo Clinic Proceedings*, **70**, 628-633.

- Sakaguchi, S., Yamaguchi, T., Nomura, T. & Ono, M. (2008). Regulatory T cells and immune tolerance. *Cell*, **133**, 775-787.
- Sayama, K., Shiraishi, S., Shirakata, Y., Kobayashi, Y., Okada, N., Okada, H. & Miki, Y. (1990). Characterization of homologous restriction factor (HRF20) in human skin and leukocytes. *Clinical & Experimental Immunology*, **82**, 355-358.
- Sayama, K., Shiraishi, S., Shirakata, Y., Kobayashi, Y. & Miki, Y. (1991a). Characterization of Decay-Accelerating Factor (DAF) in human skin. *Journal of Investigative Dermatology*, **96**, 61-64.
- Sayama, K., Shiraishi, S., Shirakata, Y., Kobayashi, Y., Seya, T. & Miki, Y. (1991b). Expression and characterization of Membrane Cofactor Protein (MCP) in human skin. *Journal of Investigative Dermatology*, **97**, 722-724.
- Schluter, H. (2000). Reversed-phase chromatography. *In Protein Liquid Chromatography*. 61. Edited by Kastner, M. Elsevier, Amsterdam. 147-223.
- Schmidt-Ullrich, R. & Paus, R. (2005). Molecular principles of hair follicle induction and morphogenesis. *Bioessays*, **27**, 247-261.
- Schweizer, J., Bowden, P. E., Coulombe, P. A., Langbein, L., Lane, E. B., Magin, T. M., Maltais, L., Omary, M. B., Parry, D. A. D., Rogers, M. A. & Wright, M. W. (2006). New consensus nomenclature for mammalian keratins. *Journal of Cell Biology*, **174**, 169-174.

Seya, T., Turner, J. R. & Atkinson, J. P. (1986). Purification and characterization of a membrane-protein (gp45-70) that is a cofactor for cleavage of C3b and C4b. *Journal of Experimental Medicine*, **163**, 837-855.

Shapiro, A. L., Vinuela, E. & Maizel, J. V., Jr. (1967). Molecular weight estimation of polypeptide chains by electrophoresis in SDS-polyacrylamide gels. *Biochemical and Biophysical Research Communications*, **28**, 815-820.

Shapiro, J. & Madani, S. (1999). Alopecia areata: diagnosis and management. *International Journal of Dermatology*, **38**, 19-24.

Siebenhaar, F., Sharov, A. A., Peters, E. M. J., Sharova, T. Y., Syska, W., Mardaryev, A. N., Freyschmidt-Paul, P., Sundberg, J. P., Maurer, M. & Botchkarev, V. A. (2007). Substance P as an immunomodulatory neuropeptide in a mouse model for autoimmune hair loss (Alopecia Areata). *Journal of Investigative Dermatology*, **127**, 1489-1497.

Silver, A. F. & Chase, H. B. (1970). DNA synthesis in the adult hair germ during dormancy (telogen) and activation (early anagen). *Developmental Biology*, **21**, 440-451.

Sinclair, R., Jolley, D., Mallari, R., Magee, J., Tosti, A., Piracinni, B. M., Vincenzi, C., Happle, R., Ferrando, J., Grimalt, R., Therese, L., Van Neste, D., Zlotogorski, A., Christiano, A. M. & Whiting, D. A. (2003). Morphological approach to hair disorders. *Journal of Investigative Dermatology Symposium Proceedings*, **8**, 56-64.

- Skinner, R. B., Light, W. H., Bale, G. F., Rosenberg, E. W. & Leonardi, C. (1995a). Alopecia areata and presence of cytomegalovirus DNA. *Jama: Journal of the American Medical Association*, **273**, 1419-1420.
- Skinner, R. B., Jr., Light, W. H., Leonardi, C., Bale, G. F. & Rosenberg, E. W. (1995b). A molecular approach to alopecia areata. *Journal of Investigative Dermatology*, **104**, S3-S4.
- Slominski, A., Wortsman, J., Plonka, P. M., Schallreuter, K. U., Paus, R. & Tobin, D. J. (2005). Hair follicle pigmentation. *Journal of Investigative Dermatology*, **124**, 13-21.
- Smith, J. C., Lambert, J. P., Elisma, F. & Figeys, D. (2007). Proteomics in 2005/2006: Developments, applications and challenges. *Analytical Chemistry*, **79**, 4325-4343.
- Smith, M. A. & Kerr, M. A. (1985). Cleavage of the second component of complement by plasma proteases: implications in hereditary C1-inhibitor deficiency. *Immunology*, **56**, 561-570.
- Starger, J. M., Brown, W. E., Goldman, A. E. & Goldman, R. D. (1978). Biochemical and immunological analysis of rapidly purified 10-nm filaments from baby hamster kidney (BHK-21) cells. *Journal of Cell Biology*, **78**, 93-109.
- Stark, H. J., Breitkreutz, D., Limat, A., Bowden, P. & Fusenig, N. E. (1987). Keratins of the human hair follicle - hyperproliferative keratins consistently expressed in outer root sheath-cells in vivo and in vitro. *Differentiation*, **35**, 236-248.

- Steinert, P. M., Jones, J. C. R. & Goldman, R. D. (1984). Intermediate filaments. *Journal of Cell Biology*, **99**, S22-S27.
- Steinert, P. M. & Parry, D. A. D. (1985). Intermediate filaments - conformity and diversity of expression and structure. *Annual Review of Cell Biology*, **1**, 41-65.
- Steinert, P. M., Parry, D. A. D. & Marekov, L. N. (2003). Trichohyalin mechanically strengthens the hair follicle - Multiple cross-bridging roles in the inner root sheath. *Journal of Biological Chemistry*, **278**, 41409-41419.
- Stenn, K. S. & Paus, R. (2001). Controls of hair follicle cycling. *Physiological Reviews*, **81**, 449-494.
- Steward, M. W. (1984). *Antibodies: Their Structure and Function*. eds. Chapman and Hall, London.
- Stout, R. D. & Bottomly, K. (1989). Antigen specific activation of effector macrophages by IFN-gamma producing (Th1) T cell clones - failure of IL-4-producing (Th2) T cell clones to activate effector function in macrophages. *Journal of Immunology*, **142**, 760-765.
- Straile, W. E., Chase, H. B. & Arsenault, C. (1961). Growth and differentiation of hair follicles between periods of activity and quiescence. *Journal of Experimental Zoology*, **148**, 205-221.
- Suckau, D., Hahner, S., Main, L. C. & Jabs, W. (2006). Application Notes # MT-83 - Accurate Quantification with ICPL: The Quantification Study of the ABRF Proteomics Research Group 2006. Bruker Daltonics GmbH.

- Sun, T. T. & Green, H. (1978). Keratin filaments of cultured human epidermal cells. Formation of intermolecular disulfide bonds during terminal differentiation. *Journal of Biological Chemistry*, **253**, 2053-2060.
- Sun, T. T., Eichner, R., Schermer, A., Cooper, D., Nelson, W. G. & Weiss, R. A. (1984). Classification, expression, and possible mechanisms of evolution of mammalian epithelial keratins: A unifying model. *In Cancer Cells. Vol. 1*. Edited by Levine, A. J., Vande Wonde, G. F., Topp, W. C. & Watson, J. D. Cold Spring Harbor Laboratory, New York. 169-176.
- Sun, T. T., Tseng, S. C. G., Huang, A. J. W., Cooper, D., Schermer, A., Lynch, M. H., Weiss, R. & Eichner, R. (1985). Monoclonal antibody studies of mammalian epithelial keratins - a review. *Annals of the New York Academy of Sciences*, **455**, 307-329.
- Sundberg, J. P., Cordy, W. R. & King, L. E. (1994). Alopecia areata in aging C3H/HeJ Mice. *Journal of Investigative Dermatology*, **102**, 847-856.
- Svanborg, C., Godaly, G. & Hedlund, M. (1999). Cytokine responses during mucosal infections: role in disease pathogenesis and host defence. *Current Opinion in Microbiology*, **2**, 99-105.
- Svejgaard, A. (2008). The immunogenetics of multiple sclerosis. *Immunogenetics*, **60**, 275-286.
- Swain, S. L., Croft, M., Dubey, C., Haynes, L., Rogers, P., Zhang, X. H. & Bradley, L. M. (1996). From naive to memory T cells. *Immunological Reviews*, **150**, 143-167.

- Switzer, R. C., 3rd, Merrill, C. R. & Shifrin, S. (1979). A highly sensitive silver stain for detecting proteins and peptides in polyacrylamide gels. *Analytical Biochemistry*, **98**, 231-237.
- Szabo, G. (1958). A regional frequency and distribution of hair follicles in human skin. *In The Biology of Hair Growth*. Edited by Montagna, W. & Ellis, R. A. Academic Press Inc., New York.
- Takata, Y., Kinoshita, T., Kozono, H., Takeda, J., Tanaka, E., Hong, K. & Inoue, K. (1987). Covalent association of C3b with C4b within C5 convertase of the classical complement pathway. *Journal of Experimental Medicine*, **165**, 1494-1507.
- Tarlow, J. K., Clay, F. E., Cork, M. J., Blakemore, A. I. F., McDonagh, A. J. G., Messenger, A. G. & Duff, G. W. (1994). Severity of alopecia areata is associated with a polymorphism in the interleukin-1 receptor antagonist gene. *Journal of Investigative Dermatology*, **103**, 387-390.
- Tazi-Ahnini, R., Cox, A., McDonagh, A. J. G., Nicklin, M. J. H., di Giovine, F. S., Timms, J. M., Messenger, A. G., Dimitropoulou, P., Duff, G. W. & Cork, M. J. (2002). Genetic analysis of the interleukin-1 receptor antagonist and its homologue IL-1L1 in alopecia areata: strong severity association and possible gene interaction. *European Journal of Immunogenetics*, **29**, 25-30.
- Tazi-Ahnini, R., Cork, M. J., Wengraf, D., Wilson, A. G., Gawkrödger, D. J., Birch, M. P., Messenger, A. G. & McDonagh, A. J. G. (2003). Notch4, a non-HLA gene in the MHC is strongly associated with the most severe form of alopecia areata. *Human Genetics*, **112**, 400-403.

- Telemo, E. & Hanson, L. A. (1996). Antibodies in milk. *Journal of Mammary Gland Biology & Neoplasia*, **1**, 243-249.
- Tenner, A. J. (1999). Membrane receptors for soluble defense collagens. *Current Opinion in Immunology*, **11**, 34-41.
- Thiel, S., Vorup-Jensen, T., Stover, C. M., Schwaeble, S., Laursen, S. B., Poulsen, K., Willist, A. C., Eggletont, P., Hansen, S., Holmskov, U., Reid, K. B. M. & Jensenius, J. C. (1997). A second serine protease associated with mannan-binding lectin that activates complement. *Nature*, **386**, 506-510.
- Tobin, D. J., Fenton, D. A. & Kendall, M. D. (1990a). Ultrastructural study of exclamation-mark hair shafts in alopecia areata. *Journal of Cutaneous Pathology*, **17**, 348-354.
- Tobin, D. J., Fenton, D. A. & Kendall, M. D. (1990b). Ultrastructural observations on the hair bulb melanocytes and melanosomes in acute alopecia areata. *Journal of Investigative Dermatology*, **94**, 803-807.
- Tobin, D. J., Orentreich, N. & Bystryn, J. C. (1994a). Autoantibodies to hair follicles in normal individuals. *Archives of Dermatology*, **130**, 395-396.
- Tobin, D. J., Orentreich, N., Fenton, D. A. & Bystryn, J. C. (1994b). Antibodies to hair follicles in alopecia areata. *Journal of Investigative Dermatology*, **102**, 721-724.
- Tobin, D. J. & Bystryn, J. C. (1996). Alopecia areata is associated with antibodies to hair follicle-specific antigens located predominantly in the

proliferative region of hair follicles. In *Hair Research for the Next Millenium*. van Neste, D. & Randall, V. A. Elsevier Science B. V., Amsterdam. 237-241.

Tobin, D. J., Hann, S. K., Song, M. S. & Bystry, J. C. (1997a). Hair follicle structures targeted by antibodies in patients with alopecia areata. *Archives of Dermatology*, **133**, 57-61.

Tobin, D. J., Sundberg, J. P., King, L. E., Boggess, D. & Bystry, J. C. (1997b). Autoantibodies to hair follicles in C3H/HeJ mice with alopecia areata-like hair loss. *Journal of Investigative Dermatology*, **109**, 329-333.

Tobin, D. J., Alhaidari, Z. & Olivry, T. (1998a). Equine alopecia areata autoantibodies target multiple hair follicle antigens and may alter hair growth. A preliminary study. *Experimental Dermatology*, **7**, 289-297.

Tobin, D. J., Olivry, T. & Bystry, J.-C. (1998b). Anti-trichohyalin antibodies in canine alopecia areata. In *Advances in Veterinary Dermatology*. Vol. 3. Edited by Kwochka, K. W., Willemse, T. & von Tscharner, C. Butterworth-Heinemann, Oxford. 355-362.

Tobin, D. J., Slominski, A., Botchkarev, V. & Paus, R. (1999). The fate of hair follicle melanocytes during the hair growth cycle. *Journal of Investigative Dermatology Symposium Proceedings*, **4**, 323-332.

Tobin, D. J. & Bystry, J. C. (2000). Immunology of alopecia areata. In *Hair and Hair Disorders: Research, Patholgy and Management*. Edited by Camacho, F. M., Randall, V. A. & Price, V. Martin Dunitz, London.

- Tobin, D. J. (2003). Plasticity and cytokinetic dynamics of the hair follicle mesenchyme during the hair growth cycle: Implications for growth control and hair follicle transformations. *Journal of Investigative Dermatology Symposium Proceedings*, **8**, 80-86.
- Tobin, D. J., Gardner, S. H., Luther, P. B., Dunston, S. M., Lindsey, N. J. & Olivry, T. (2003). A natural canine homologue of alopecia areata in humans. *British Journal of Dermatology*, **149**, 938-950.
- Tobin, D. J. & Olivry, T. (2004). Spontaneous Canine Model of Alopecia Areata. In *Animal models of human inflammatory skin diseases*. Edited by Chan, L. S. CRC Press, US. 469-481.
- Tobin, D. J. (2005). The Biogenesis and Growth of Human Hair. In *Hair in Toxicology: An Important Bio-Monitor*. Tobin, D. J. Royal Society of Chemistry, Cambridge. 3-33.
- Tobin, D. J. (2006). Biochemistry of human skin - our brain on the outside. *Chemical Society Reviews*, **35**, 52-67.
- Tonegawa, S. (1983). Somatic generation of antibody diversity. *Nature*, **302**, 575-581.
- Tosti, A., LaPlaca, M., Placucci, F., Gentilomi, G., Venturoli, S., Zerbini, M. & Musiani, M. (1996). No correlation between cytomegalovirus and alopecia areata. *Journal of Investigative Dermatology*, **107**, 443-443.
- Towbin, H., Staehelin, T. & Gordon, J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure

and some applications. *Proceedings of the National Academy of Sciences of the United States of America*, **76**, 4350-4354.

Trichohyalin (N-16): sc-47517 Specifications. Santa Cruz Biotechnology Inc., California.

Trouw, L. A., Blom, A. M. & Gasque, P. (2008). Role of complement and complement regulators in the removal of apoptotic cells. *Molecular Immunology*, **45**, 1199-1207.

Tseng, S. C. G., Jarvinen, M. J., Nelson, W. G., Huang, J. W., Woodcock-Mitchell, J. & Sun, T. T. (1982). Correlation of specific keratins with different types of epithelial differentiation - monoclonal antibody studies. *Cell*, **30**, 361-372.

Tupin, E., Kinjo, Y. & Kronenberg, M. (2007). The unique role of natural killer T cells in the response to microorganisms. *Nature Reviews Microbiology*, **5**, 405-417.

Tyers, M. & Mann, M. (2003). From genomics to proteomics. *Nature*, **422**, 193-197.

Uchi, H., Terao, H., Koga, T. & Furue, M. (2000). Cytokines and chemokines in the epidermis. *Journal of Dermatological Science*, **24**, S29-S38.

van Eyk, J. E. (2001). Proteomics: Unraveling the complexity of heart disease and striving to change cardiology. *Current Opinion in Molecular Therapeutics*, **3**, 546-553.

- van Neste, D. & Tobin, D. J. (2004). Hair cycle and hair pigmentation: dynamic interactions and changes associated with aging. *Micron*, **35**, 193-200.
- van Scott, E. J. (1958). Morphologic changes in pilosebaceous units and anagen hairs in alopecia areata. *Journal of Investigative Dermatology*, **31**, 35-43.
- Villadangos, J. A. & Schnorrer, P. (2007). Intrinsic and cooperative antigen-presenting functions of dendritic-cell subsets in vivo. *Nature Reviews Immunology*, **7**, 543-555.
- Vorup-Jensen, T., Petersen, S. V., Hansen, A. G., Poulsen, K., Schwaeble, W., Sim, R. B., Reid, K. B. M., Davis, S. J., Thiel, S. & Jensenius, J. C. (2000). Distinct pathways of mannan-binding lectin (MBL)- and C1-complex autoactivation revealed by reconstitution of MBL with recombinant MBL-associated serine protease-2. *Journal of Immunology*, **165**, 2093-2100.
- Wallevik, K. (1973). Reversible denaturation of human serum albumin by pH, temperature, and guanidine hydrochloride followed by optical rotation. *Journal of Biological Chemistry*, **248**, 2650-2655.
- Walport, M. J. (2001). Advances in immunology: Complement (First of two parts). *New England Journal of Medicine*, **344**, 1058-1066.
- Webermatthiesen, K. & Sterry, W. (1990). Organization of the Monocyte Macrophage System of Normal Human Skin. *Journal of Investigative Dermatology*, **95**, 83-89.

- Weiler, J. M., Daha, M. R., Austen, K. F. & Fearon, D. T. (1976). Control of amplification convertase of complement by plasma protein beta-1h. *Proceedings of the National Academy of Sciences of the United States of America*, **73**, 3268-3272.
- Welsh, E. A., Clark, H. H., Epstein, S. Z., Reveille, J. D. & Duvic, M. (1994). Human leukocyte antigen-DQB1*03 alleles are associated with alopecia areata. *Journal of Investigative Dermatology*, **103**, 758-763.
- Wengraf, D. A., McDonagh, A. J. G., Lovewell, T. R. J., Vasilopoulos, Y., Macdonald-Hull, S. P., Cork, M. J., Messenger, A. G. & Tazi-Ahnini, R. (2008). Genetic analysis of autoimmune regulator haplotypes in alopecia areata. *Tissue Antigens*, **71**, 206-212.
- Westgate, G. E., Gibson, W. T., Kealey, T. & Philpott, M. P. (1993). Prolonged Maintenance of Human Hair-Follicles in-Vitro in a Serum-Free Medium. *British Journal of Dermatology*, **129**, 372-379.
- Whiting, D. A. (2004). The structure of the human hair follicle - light microscopy of vertical and horizontal sections of scalp biopsies. eds. Canfield Publishing, Fairfield.
- Winter, H., Langbein, L., Praetzel, S., Jacobs, M., Rogers, M. A., Leigh, I. M., Tidman, N. & Schweizer, J. (1998). A novel human type II cytokeratin, K6hf, specifically expressed in the companion layer of the hair follicle. *Journal of Investigative Dermatology*, **111**, 955-962.
- Woof, J. M. & Kerr, M. A. (2006). The function of immunoglobulin A in immunity. *Journal of Pathology*, **208**, 270-282.

- Xiao, F. L., Yang, S., Lin, G. S., Gao, M., Cui, Y., Yin, X. Y., Wang, P. G., Xu, S. X. & Zhang, X. J. (2007). HLA haplotypic association with different phenotype of alopecia areata in Chinese Hans. *Journal of Dermatological Science*, **45**, 206-209.
- Zillikens, D., Rose, P. A., Balding, S. D., Liu, Z., Olague-Marchan, M., Diaz, L. A. & Giudice, G. J. (1997). Tight clustering of extracellular BP180 epitopes recognized by bullous pemphigoid autoantibodies. *Journal of Investigative Dermatology*, **109**, 573-579.
- Zoller, M., McElwee, K. J., Engel, P. & Hoffmann, R. (2002). Transient CD44 variant isoform expression and reduction in CD4⁺/CD25⁺ regulatory T cells in C3H/HeJ mice with alopecia areata. *Journal of Investigative Dermatology*, **118**, 983-992.