

Joseph O. WARHAM

MAPPING BIOSPHERE STRONTIUM ISOTOPE RATIOS ACROSS MAJOR LITHOLOGICAL BOUNDARIES

Keywords: Archaeology, geochemistry, precipitation, residential mobility, Sr-isotopes, stream water, TIMS, vegetation

ABSTRACT

Strontium isotope analysis has provided archaeologists with an unprecedented opportunity to study the mobility of humans and animals in the past. However, a lack of systematic environmental baseline data has seriously restricted the full potential of the analytical technique; there is little biosphere data available against which to compare measured skeletal data.

This thesis examines the extent to which geographic variation in biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ composition can be spatially resolved within the lowland terrain of England, in a geographically and geologically coherent study area. Systematically collected samples of vegetation, stream water and surface soils, including new and archived material have been used. The potential of these sample media to provide reliable estimates of the $^{87}\text{Sr}/^{86}\text{Sr}$ composition of bioavailable strontium are evaluated under both high-density and low-density sampling regimes, and against new analyses of local archaeological material.

Areas lying south of the Anglian glacial limit, display a pattern of geographic $^{87}\text{Sr}/^{86}\text{Sr}$ biosphere variation (0.7080–0.7105) controlled by solid geology, as demonstrated by high-density biosphere mapping. Data collected at a wider geographic scale, including above superficial deposits, indicate the dominant influence of re-worked local rocks on the biosphere. These methods have enabled a reclassification of the archaeologically important Cretaceous Chalk domain. Analysis of rainwater and other indicators of atmospheric deposition show that, in this setting, local biosphere variation is not significantly perturbed by atmospheric inputs.

Time-related data from archaeological cattle and sheep/goat tooth enamel suggest that the modern biosphere data can be used to understand livestock management regimes and that these are more powerful than using an average value from the enamel. A more complete understanding of possible patterns of mobility in a group of humans has been achieved through analysis of material from Winchester and comparison with the Chalk biosphere domain.

ACKNOWLEDGEMENTS

I would like to thank the British Geological Survey's British University Funding Initiative (BUFI) and the School of Life Sciences at the University of Bradford for jointly funding this project; I am sincerely grateful to the Division of Archaeological, Geographical and Environmental Sciences (AGES) for awarding me the studentship and giving me the opportunity to pursue doctoral research. It is a great pleasure also to acknowledge the contributions of my supervisors, Dr Janet Montgomery (Durham University, Department of Archaeology), Professor Jane Evans (NIGL), Dr Louise Ander (BGS), Dr David Cotton (AGES), and latterly, Dr Cathy Batt (AGES) and Professor Carl Heron (AGES). It has been a privilege to benefit from their knowledge and experience, their astute guidance, insight and patience.

Fundamentally, this research project could not have been undertaken without access to facilities provided by the NERC Isotope Geosciences Laboratory (NIGL), and the support of the scientists who have made me so welcome at the British Geological Survey (BGS). Accordingly, I would like to thank the Head of NIGL, Professor Randall Parish, for accepting an archaeologist into the fold and, amongst others, Dr Nicola Atkinson, Neil Boulton, Carolyn Chenery, Dr Simon Chenery, Dr Ian Millar, Dr Steve Noble, Aaran Sumner, and Vanessa Pashley. I should also like to thank my office-mates Dr Anglea Lamb and Dr Andrea Snelling, for their constant encouragement during the final phases of the production of this PhD thesis.

This research was also dependent to a great extent on the availability of archived materials, including both environmental samples and archaeological finds. The maintenance of such archives represents a significant commitment on the part of the institutions and individuals responsible for them, and an important investment for future research. Thus, the Geochemical Baselines and Medical Geology group at the BGS made a substantial contribution to the success of this project. I am grateful to Kate Knights, and Andrea Mills for their assistance in accessing material held within the G-BASE sample archive (Chapters 4 and 5), and also to Dr Andreas Scheib for the training and supervision he provided during the GEMAS geochemical mapping project (Chapter 5). Heath Malcolm and Hayley Guyatt, of the NERC Centre for Ecology and Hydrology (CEH), provided me with access to unpublished data and to samples from the Rural Heavy Metals Monitoring Network, as well as advice on rainwater sampling without which the work presented in Chapter 6 of this thesis would not have been possible.

Thanks are also due to the curators who gave me permission to carry out the destructive analysis of archaeological material (Chapters 7 and 8), as follows: Dr Esther Cameron, Curator of Archaeology for Oxfordshire Museums Service; Brett Thorn, Curator of Archaeology for the Buckinghamshire County Museum; and Helen Rees, Curator of Archaeology for Winchester Museums. I am indebted also to Dr Annia Cerryson and to Dr Jo Buckberry (AGES) for their insights into Early Medieval burial practices, and for providing pre-publication access to their work on the human skeletal material from the site of Old Dairy Cottage (Chapter 8). I would also like to thank James Rackham (The Environmental Archaeology Consultancy), Dr Rebecca Nicholson (Oxford Archaeology) and Dr Fay Worley (English Heritage) for their advice on archaeozoology.

I owe a debt of thanks to all of my colleagues, academic peers, and friends whose own research, comments and questions have encouraged, challenged and enlightened me in equal measure. My deepest gratitude goes to my family, especially Gemma Martin and my parents, Richard and Ilona Warham, to my brother Peter, and Nana Dorothy; they have provided steadfast encouragement and support throughout my mounting years of study, and at times some much-needed perspective on life beyond the lab, library and laptop.

CONTENTS

LIST OF FIGURES	vii
LIST OF TABLES	x
CHAPTER 1	
INTRODUCTION	1
1.1 Aims and objectives	4
1.2 Thesis structure	6
CHAPTER 2	
LITERATURE REVIEW	8
2.1 The characterisation of biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ values for archaeological applications	12
2.2 Systematic geochemical mapping within the British Isles	22
2.4 Sample media available for biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ mapping	29
CHAPTER 3	
MATERIALS AND METHODS	36
3.1 Analytical method	36
3.1.1 Cation exchange chemistry	37
3.1.2 TIMS analysis of $^{87}\text{Sr}/^{86}\text{Sr}$	38
3.1.3 Isotope dilution	40
3.2 Modern environmental sample media	41
3.2.1 Vegetation (Wood, Leaf, and Grass samples)	41
3.2.1a Disaggregation and homogenisation	42
3.2.1b Microwave-assisted extraction of strontium	43
3.2.1c Blank characteristics	44
3.2.1d Procedural reproducibility	45
3.2.2 Waters (Water and Rain samples)	48
3.2.2a Water sample treatment	49
3.2.2b Stream water (Water samples)	50
3.2.2c Rainwater (Rain samples)	51
3.2.3 Surface soil samples (Soil samples)	51
3.2.3a Soil-leach method	52
3.2.3b Blank characteristics	53
3.3 Archaeological sample media	54
3.3.1 Faunal material	55
3.3.1a Sampling regime used to allow $^{87}\text{Sr}/^{86}\text{Sr}$ variation in sheep and cattle teeth to be plotted on common time-related axis	56
3.3.1b Hypsodont tooth sampling methods	61
3.3.1c Comparison of bulk samples and weighted means	62
3.3.1d Duplicate analyses	64
3.3.2 Human material	67
3.3.3 Preparation of dental tissues for analysis	68
3.3.4 Blank characteristics	69

CHAPTER 4		
	HIGH DENSITY BIOSPHERE ⁸⁷Sr/⁸⁶Sr MAPPING	70
4.1	Characterisation of local conditions	71
4.1.1	Geology	71
4.1.2	Surface hydrology	75
4.1.3	Soils and modern land-use	75
4.2	Biosphere ⁸⁷ Sr/ ⁸⁶ Sr characterisation	77
4.2.1	Results	78
4.2.2	Discussion	81
4.2.3	Conclusions	84
4.3	Variation in ⁸⁷ Sr/ ⁸⁶ Sr composition within geological divisions of the biosphere	86
4.3.1	Variation with distance from Chalk escarpment	87
4.3.1a	Discussion	90
4.3.2	Variation above the Oxford Clay Formation	91
4.3.2a	Discussion	93
4.3.3	Conclusions	95
CHAPTER 5		
	REFINING OUR UNDERSTANDING OF THE INFLUENCE OF SUPERFICIAL DEPOSITS ON LARGE-SCALE GEOGRAPHIC BIOSPHERE ⁸⁷Sr/⁸⁶Sr VARIATION	97
5.1	The use of a low density survey to overcome the geographic focus of detailed studies	98
5.1.1	GEMAS sample site selection and field methodology	99
5.1.2	Results	102
5.1.3	Discussion	103
5.1.4	Conclusions	106
5.2	The influence of glacial till above the Chalk	107
5.2.1	Sample Selection	108
5.2.2	Results	110
5.2.3	Discussion	111
5.2.4	Conclusions	114
5.3	The Clay-with-flints: the influence of decalcified in-situ sediments on biosphere variation above the Chalk	114
5.3.1	Geochemical reconnaissance above the Clay-with-flints	116
5.3.2	Biosphere sampling exercise targeting Clay-with-flints	118
5.3.3	Results	120
5.3.4	Discussion	120
5.3.5	Conclusions	123
5.4	The coastal wetlands: a distinctive biosphere domain?	124
5.4.1	The Fenland Basin of eastern England	126
5.4.2	Sample Selection	127
5.4.3	Results	129
5.4.4	Discussion	129
5.4.5	Conclusions	132

CHAPTER 6		
	CHARACTERISING THE INFLUENCE OF ATMOSPHERIC INPUT	133
6.1	The immediate impact of volcanic ash originating from the Eyjafjallajökull eruption (Iceland) in 2010	134
6.1.1	Sample collection	135
6.1.2	Results	137
6.1.3	Discussion	139
6.1.4	Conclusions	141
6.2	Comparison of the isotopic composition of woody material and accompanying foliage	142
6.2.1	Results	143
6.2.2	Discussion	145
6.2.3	Conclusions	148
6.3	Evidence regional variation in the $^{87}\text{Sr}/^{86}\text{Sr}$ composition of atmospheric inputs within the UK	148
6.3.1	Results	150
6.3.2	Discussion	154
6.3.3	Conclusions	161
CHAPTER 7		
	AN EVALUATION OF THE POTENTIAL OF BIOSPHERE MAPPING IN THE CONTEXT OF FAUNAL DATA FROM ARCHAEOLOGICAL SITES	163
7.1	The identification of suitable archaeological faunal assemblages: opportunities and constraints	164
7.1.1	Ivinghoe Beacon and Ward's Coombe: two sites above the Chalk <i>sensu lato</i>	168
7.1.2	Bierton Vicarage Garden: a site located above the Upper Jurassic mid-vale ridge	171
7.1.3	Bicester Fields Farm: a site between the Oxford Clay and the Great Oolite	173
7.2	Characterisation of the age-related bias in the selected faunal material	174
7.3	Assessment of diagenesis	176
7.4	Results	180
7.4.1	Ivinghoe Beacon and Ward's Coombe	184
7.4.2	Bierton Vicarage Garden	186
7.4.3	Bicester Fields Farm	187
7.5	Discussion	189
7.6	Conclusions	195
CHAPTER 8		
	AN EVALUATION OF CONSTRAINTS ON THE INTERPRETATION OF HUMAN $^{87}\text{Sr}/^{86}\text{Sr}$ DATA	197
8.1	The significance for strontium isotope studies of the geographic distribution of putative Anglo-Saxon execution cemeteries	198
8.1.1	A human $^{87}\text{Sr}/^{86}\text{Sr}$ range defined for southern England?	201
8.1.2	Site description	212
8.1.3	Historical background	214
8.2	Description of samples	216
8.3	Assessment of diagenesis	216
8.4	Results	218
8.5	Discussion	219
8.6	Conclusions	222

CHAPTER 9		
	CONCLUSIONS	224
9.1	Findings summarised with reference to original aims	226
9.2	Limitations	228
9.3	Recommendations for techniques of biosphere mapping	229
9.4	Implications for future research	232
REFERENCES		236
PERSONAL COMMUNICATIONS		256
APPENDICES		
APPENDIX A	METHODS	
APPENDIX B	ARCHAEOLOGICAL MATERIALS	
APPENDIX C	DATA TABLES	

LIST OF FIGURES

CHAPTER 2

- 2.01 Bar chart indicating the rapid increase in publications relating to the use of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis within archaeology, anthropology and evolutionary biology, derived from Web of Science data-base query (Web of Knowledge 2011). 9
- 2.02 Schematic diagram showing the delineation of a drainage basin by watershed and the 1x1 km Ordnance Survey (OS) grid-based system used by the G BASE project to achieve consistent sample density, after Johnson *et al.* (2008b: Figure 4.1) with modifications. 27

CHAPTER 3

- 3.01 Illustration showing the macroscopic sampling method uses to obtain serial enamel samples removed from cattle and sheep teeth. 61
- 3.02 Scatter plot comparing cattle and sheep/goat teeth from Bicester Fields Farm on the basis of bulk enamel (■), and weighted mean enamel (□) $^{87}\text{Sr}/^{86}\text{Sr}$ composition and Sr concentration (cf. Tables 3.13 and 3.14). 64
- 3.03 Scatter plot comparing principal series of enamel samples A, B, and C (□) with replicate series A2 and C2 (■) from cattle tooth BIFF_04_M2 on the basis of: A) $^{87}\text{Sr}/^{86}\text{Sr}$ composition; B) strontium concentration (cf. Table 3.13). 65

CHAPTER 4

- 4.01 Location of field area, showing all G-BASE drainage sample sites above a generalised map of bedrock geology and topographic relief. 71
- 4.02 Simplified SE–NW sketch section across study area, with exaggerated y-axis (not to scale). 72
- 4.03 Map showing location of samples selected for analysis from first and second order catchments (Wood and Water) and surface soil samples (Soil) in relation to bedrock geology. 78
- 4.04 Individual value plot of Wood, Water and Soil $^{87}\text{Sr}/^{86}\text{Sr}$ data for samples collected across the principal lithologies within the study area. 82
- 4.05 Map showing sample sites selected to investigate the effect of catchment size and complexity in relation to underlying geology. 87
- 4.06 Line chart showing changes in the $^{87}\text{Sr}/^{86}\text{Sr}$ composition of Soil, Wood and Water at an increasing distance from the foot of the Chalk escarpment, in relation to topography and bedrock geology (cf. Figure 4.06). 89
- 4.07 Individual value plot showing proposed biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ ranges (mean \pm 2 SD) based on Wood samples associated with geological divisions of the biosphere that are contiguous with the Peterborough member of the Oxford Clay formation. 95

CHAPTER 5

- 5.01 Map showing location of low-density GEMAS Grass samples selected for $^{87}\text{Sr}/^{86}\text{Sr}$ analysis, in relation to generalised map of underlying bedrock geology. 102
- 5.02 Individual value plot comparing low-density data from GEMAS (■) with previously published low-density vegetation $^{87}\text{Sr}/^{86}\text{Sr}$ data from samples collected in similar geological settings. 104

5.03	Map showing low-density distribution of selected stream water samples (G-BASE) in relation to glacial till and generalised map of underlying bedrock.	109
5.04	Individual value plot comparing the $^{87}\text{Sr}/^{86}\text{Sr}$ composition of G-BASE stream water samples (Water) collected from areas underlain by glacial till, with previously established ranges associated with Chalk bedrock geology.	113
5.05	Map showing Sr concentration in G-BASE surface soils that are located above outcrops belonging to the Chalk Group in relation to topography.	117
5.06	Map showing Sr concentration in G-BASE surface soils that are located above outcrops belonging to the Chalk Group in relation to superficial geology.	117
5.07	Map showing location of G-BASE surface soil samples (Soil), and vegetation (Wood) samples collected in order to assess the influence of the Clay-with-flints on the Chalk biosphere <i>sensu lato</i> , and the attenuation of the marine carbonate influence beyond the Chalk escarpment of Buckinghamshire, in relation to underlying geology.	118
5.08	Line chart showing changes in $^{87}\text{Sr}/^{86}\text{Sr}$ composition of soil leaches (Soil) and vegetation (Wood) in relation to topography, underlying geology and the range of values previously obtained locally from samples collected above the Lower Cretaceous mudrocks (Gault).	121
5.09	Map showing the location of G-BASE stream water sampling sites selected to investigate biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ variation within the Fenland Basin of eastern England. Sites are shown in relation to underlying superficial deposits and the proximal GEMAS grass samples (cf. Figure 5.01).	128
5.10	Individual value plot comparing G-BASE stream water $^{87}\text{Sr}/^{86}\text{Sr}$ data (Water) from the Fenland Basin with data from surrounding regions and proximal GEMAS Grass samples.	130

CHAPTER 6

6.01	Line and bar chart showing the results of the $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of Grass and Rain samples collected from 25/04/2010–12/05/2010 (documented in Appendix C).	138
6.02	Scatter plot showing the linear regression (ordinary least squares) of the $^{87}\text{Sr}/^{86}\text{Sr}$ composition of G-BASE Wood versus Leaf samples.	144
6.03	Maps showing the location of the rainwater collection sites comprising the CEH Rural Heavy Metals Monitoring Network and the accompanying $^{87}\text{Sr}/^{86}\text{Sr}$ values reported in Table 6.04, together with the published rainwater data from the UK.	149
6.04	Histograms of annual monthly rainwater $^{87}\text{Sr}/^{86}\text{Sr}$ composition formulated using deposition weightings (A) and frequency distribution (B).	153
6.05	Graph showing the relationship between rainwater $^{87}\text{Sr}/^{86}\text{Sr}$ composition and $1/\text{Sr}$.	157
6.06	Graph showing the relationship between rainwater $^{87}\text{Sr}/^{86}\text{Sr}$ composition and enrichment in non-marine cations (Ba and Al).	157

CHAPTER 7

7.01	Map showing the location of selected Iron Age archaeological sites in Oxfordshire and Buckinghamshire, shown in relation to distribution of superficial (glacial till, river terraces and alluvium) and bedrock geology.	167
------	--	-----

7.02	Scatter plot of $^{87}\text{Sr}/^{86}\text{Sr}$ versus Sr ($\mu\text{g/g}$), comparing sheep/goat dentine (closed symbols) with sheep/goat enamel (open symbols).	177
7.03	Hypothetical mixing lines illustrating the possible magnitude of strontium diagenesis on dentine within a Chalk burial environment, allowing only for the deposition of secondary minerals with a $^{87}\text{Sr}/^{86}\text{Sr}$ composition of 0.7075.	178
7.04	Scatter plot of Sheep/goat data from Chalk burial environments (Ward's Coombe and Ivinghoe Beacon) shown as $^{87}\text{Sr}/^{86}\text{Sr}$ versus $1/\text{Sr}$, highlighting a possible mixing line within the enamel data obtained from one individual (IB_02) recovered from the site of Ivinghoe Beacon.	180
7.05	Individual value plot comparing the concentration of strontium in modern sample media – Water < Soil (leachable) << Wood < Grass Leaf << Soil (bulk) – with that in archaeological cattle and sheep/goat dental enamel.	183
7.06	Line charts showing $^{87}\text{Sr}/^{86}\text{Sr}$ composition of serial enamel samples from (A) cattle, and (B) sheep/goat teeth obtained from Ivinghoe Beacon and Ward's Coombe, plotted on common developmental timeline.	185
7.07	Line charts showing $^{87}\text{Sr}/^{86}\text{Sr}$ composition of serial enamel samples from (A) cattle, and (B) sheep/goat teeth obtained from Bierton Vicarage Garden, plotted on common developmental timeline.	187
7.08	Line charts showing $^{87}\text{Sr}/^{86}\text{Sr}$ composition of serial enamel samples from (A) cattle, and (B) sheep/goat teeth from Bicester Fields Farm, plotted on common developmental timeline.	189

CHAPTER 8

8.01	Distribution of Anglo-Saxon execution cemeteries (No 23 = Old Dairy Cottage) identified by Reynolds (2009) in relation to Cretaceous bedrock geology dominated by the Upper Cretaceous Chalk and the Lower Cretaceous Gault Formations.	199
8.02	Summary biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ data from Hubbard Brook Experimental Forest (Blum <i>et al.</i> 2000 ; Blum <i>et al.</i> 2001), indicating the consistent central tendency within the local ecosystem and the change in variance between trophic levels.	202
8.03	Individual value plot demonstrating the narrowing of variance associated with iteratively re-sampling an underlying biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ distribution.	204
8.04	Hypothetical mixing line between end-members representing the Chalk biosphere <i>sensu lato</i> .	207
8.05	Box-plots displaying archaeological human $^{87}\text{Sr}/^{86}\text{Sr}$ data drawn from communal burial grounds in England shown in relation to potential significant break points in the predicted range for human variation above the Chalk <i>sensu lato</i> .	210
8.06	Comparison of enamel and dentine data from Sk578 to published human dentine data from other Chalk burial environments.	217
8.07	Graph showing enamel results from Old Dairy Cottage in relation to reference lines suggested in Table 8.01	220

CHAPTER 9

9.01	Map showing a generalised scheme of $^{87}\text{Sr}/^{86}\text{Sr}$ variation within the Chalk biosphere package <i>sensu lato</i> .	225
------	--	-----

LIST OF TABLES

CHAPTER 2

2.01	Characteristics of the sample media used commonly in geochemical mapping exercises.	24
2.02	Strontium concentrations in water, humus, topsoil and stream sediment reported in the “Statistical data of analytical results” provided in the Geochemical Atlas of Europe (Salminen <i>et al.</i> 2005: 99–108).	25
2.03	Potential sample media for use in the characterisation of biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ values, after Evans & Tatham (2004: Table 1) with modifications	31
2.04	Examples of soil-leach methods used to characterise the distribution and relative mobility of metals in soils.	33

CHAPTER 3

3.01	Mean values for replicate measurements of NBS 987 used to normalise $^{87}\text{Sr}/^{86}\text{Sr}$ data collected in static mode.	39
3.02	Procedural blanks for the microwave-assisted digestion of plant material (Wood, Leaf and Grass).	45
3.03	Results of the replicate $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of NIST Standard Reference Material 1515 (Apple Leaves) analysed by TIMS.	47
3.04	Results of the initial $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of vegetation samples (Wood), and replicate analyses undertaken in chemistry batch P492	48
3.05	Results of the $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of field duplicates of vegetation samples (Wood).	48
3.06	Results indicating agreement between ID-TIMS and G-BASE concentration of strontium in stream water samples (Water) from chemistry batch P455.	50
3.07	Results for G-BASE field blanks and procedural blank for preparation of stream water samples (Water) in chemistry batch P455.	50
3.08	Analysis of G-BASE stream water field duplicates.	51
3.09	Mandibular molar formation times of cattle (Brown <i>et al.</i> 1960) and sheep (Milhaud and Nézit 1991).	57
3.10	Estimated rate of formation of the enamel crowns of the second (M_2) and third (M_3) mandibular molars of cattle & sheep/goat. Formation time is quoted as a central value within the ranges cited in Table 3.09.	60
3.11	Details of bulk and serial samples of cattle enamel obtained from the same crowns.	63
3.12	Details of bulk and serial samples of sheep/goat enamel obtained from the same crowns.	64
3.13	Details of duplicate serial samples of cattle enamel from the same lobe of the second molar (M_2) of cattle mandible BIFF_04.	65
3.14	Permanent tooth formation time following Montgomery (2002: 52, Table 3.3).	67

CHAPTER 4

4.01	Local geological succession following Horton <i>et al.</i> (1995).	74
4.02	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of Wood and Water, and Soil samples from geologically comparable locations (cf. Table 4.03).	79
4.03	Key to stratigraphic column in Table 4.02.	80

4.04	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis vegetation (Wood) and stream water (Water) collected at an increasing distance from the foot of the Chalk escarpment (cf. Figure 4.05).	88
4.05	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of soil leaches (Soil) from samples collected at an increasing distance from the foot of the Chalk escarpment (cf. Figure 4.05).	88
4.06	Result of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of Wood sample collected within a simple first-order catchment location above the Peterborough member of the Oxford Clay (PET), within a watershed fed by drainage from the younger members of the Ancholme Group (WEY, SBY).	92
4.07	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of Wood samples collected within the same drainage basin above the Peterborough member (PET) of the Oxford Clay, at locations associated with increasingly complex catchment characteristics.	92
4.08	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of Wood samples collected at complex, high-order (3 rd order and higher) locations above the Stewartby (SBY) and Peterborough (PET) members of the Oxford Clay.	92

CHAPTER 5

5.01	The effect of cartographic scale on map detail and information content.	98
5.02	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of GEMAS Grass samples.	103
5.03	Results of the analysis of G-BASE stream waters obtained from areas underlain by glacial till that lies above Cretaceous and Jurassic bedrock geology.	111
5.04	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of vegetation samples (Wood) used to investigate the influence of the Clay-with-flints on the Chalk biosphere <i>sensu lato</i> in the Chalk escarpment zone (cf. Figure 5.07).	120
5.05	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of G-BASE surface soil samples (Soil) used to investigate the influence of the Clay-with-flints on the Chalk biosphere <i>sensu lato</i> in the Chalk escarpment zone (cf. Figure 5.07).	120
5.06	Summary of available biosphere data (vegetation only) that are characteristic of the Chalk biosphere <i>sensu stricto</i> .	124
5.07	Results of the analysis of G-BASE stream water samples selected to investigate biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ variation across the Fenland Basin.	129

CHAPTER 6

6.01	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of paired Wood and Leaf samples presented in ascending order of Wood composition.	143
6.02	Details of the rainwater samples from the CEH Rural Heavy Metals Monitoring Network collection site in Oxfordshire (Wytham) collected during 2008, and results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis.	151
6.03	Details of the rainwater samples from the CEH Rural Heavy Metals Monitoring Network selected for $^{87}\text{Sr}/^{86}\text{Sr}$ analysis, which provide coverage across the British Isles.	151
6.04	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of rainwater samples from the CEH Rural Heavy Metals Monitoring Network, which provide coverage across the British Isles.	152

CHAPTER 7

7.01	Simplified Iron Age chronological scheme for lowland Britain based on Cunliffe (2005: 22, Figure 2.2) and indicative date-ranges associated with the selected sites used within this study.	167
7.02	Age stage of selected cattle and sheep/goat mandibles estimated following the recommendations of Hambleton (1999) using mandible wear score allocated following Grant (1982).	175
7.03	Results of serial $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of cattle molars.	181
7.08	Results of serial $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of sheep/goat molars.	182

CHAPTER 8

8.01	Significant break-points in the biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ range associated human and faunal averages obtained from the Chalk <i>sensu lato</i> .	209
8.02	Selected burial data for articulated individuals from Old Dairy Cottage following Cherryson and Buckberry (in preparation).	214
8.03	Radiocarbon dates from burials at Old Dairy Cottage following Cherryson and Buckberry (in preparation).	214
8.04	Results of $^{87}\text{Sr}/^{86}\text{Sr}$ analysis of enamel samples from Old Dairy Cottage.	219