

List of Figures

2.1	A map showing the find sites of RLWm ware across the eastern Mediterranean.	20
2.2	Maps showing the distribution of RLWm ware (a) up to 1400 BC and (b) after 1400 BC.	21
2.3	The three main forms of RLWm ware – (a) spindle bottle, (b) pilgrim or lentoid flask, (c) arm-shaped vessel.	25
2.4	Thin section of a fine RLWm ware sherd from Saqqara, Egypt under crossed polars.	27
2.5	Thin section of a coarse RLWm ware sherd from Kilise Tepe under crossed polars.	27
2.6	The interior surface of sherd 6 from Arpera, Cyprus, showing marks created by the wheel throwing process.	29
2.7	The stages in the reconstruction of a lentoid flask by a modern potter.	31
2.8	Burnishing marks on the exterior of sherd 6 from Arpera, Cyprus.	32
2.9	Pot marks on the base of a spindle bottle from Arpera, Cyprus (sherd 1), and the handle of a pilgrim flask from Dhenia, Cyprus (sherd 1).	34
3.1	Maps showing the changes in the political systems during the Late Bronze Age.	49
4.1	Map of Cyprus showing sources of sherd material.	59
4.2	Sherd 2 from Arpera, Cyprus, showing the pot mark inscribed on the base.	62
4.3	Sherd 4 from Arpera, Cyprus, showing the pot mark inscribed on the base and the marks of the wheel throwing on the interior.	62
4.4	The exterior and interior of sherd 9 from Arpera, Cyprus.	62
4.5	Pilgrim flask sherd from Dhenia, Cyprus.	64
4.6	The interior and exterior of (a) sherd 1 and (b) sherd 2 from Enkomi, Cyprus.	67
4.7	Sherd 3 from Hala Sultan Tekke, Cyprus.	70
4.8	Sherd 6 from Hala Sultan Tekke, Cyprus.	70
4.9	Sherd 8 from Hala Sultan Tekke, Cyprus.	71
4.10	Sherd 9 and sherd 11 from Hala Sultan Tekke, Cyprus.	72
4.11	Sherd 2 from Kalavassos – <i>Ayios Dimitrios</i> , Cyprus.	76
4.12	Sherd 1 from Kalavassos – <i>Ayios Dimitrios</i> , Cyprus.	76
4.13	Sherd 9 from Kalavassos – <i>Ayios Dimitrios</i> , Cyprus.	76

4.14	The exterior and interior of sherd 7 from Kalavassos, Cyprus.	77
4.15	The exterior and interior of sherd 9 from Kalavassos, Cyprus.	77
4.16	The exterior and interior of sherd 4 from Kazaphani, Cyprus showing the similarity between the two surfaces.	79
4.17	Sherd 5 from Kazaphani, Cyprus.	80
4.18	Sherd 10 from Kazaphani, Cyprus.	80
4.19	Sherd 5 from Kouklia, Cyprus.	84
4.20	Sherd of a spindle bottle from the site of Myrtou-Pigadhes, Cyprus.	86
4.21	Map of Turkey showing sources of sherd material.	86
4.22	Sherd 34 from Boğazköy, Turkey.	90
4.23	The exterior and interior surfaces of sherd 205 from Kilise Tepe, Turkey.	95
4.24	The exterior and interior of sherd 206 from Kilise Tepe, Turkey.	95
4.25	The exterior and interior of sherd 1 from Kuşakli, Turkey.	98
4.26	Sherd 4 from Kilise Tepe, Turkey.	98
4.27	Map showing the location of the site of Tell Tweini, Syria.	99
4.28	Location of the site of Saqqara, Egypt.	101
4.29	The exterior and interior of sherd 2 from the site of Saqqara, Egypt.	103
4.30	Sherd 1 from Saqqara, Egypt, with the visible residue adhering to the interior surface.	107
5.1	Chemical reactions involved in methylation, silylation and transesterification.	116
5.2	Three sherds exhibiting a white deposit – from the left Arpera sherd 3, Arpera sherd 5 and Hala Sultan Tekke sherd 7.	125
5.3	Process of saponification of triglycerol, in this example tripalmitin, to glycerol and free palmitic acid.	127
5.4	Examples of local and RLWm ware sherds from Boğazköy.	140
7.1	Structures of wax esters and hydroxyl wax esters.	177
7.2	Chromatogram of the beeswax residue from the interior of sherd 34 from Boğazköy.	180
7.3	(a) mass spectrum of C ₂₇ <i>n</i> -alkane; (b) mass spectrum of C ₂₄ alcohol; (c) mass spectrum of C ₄₀ palmitic wax ester.	181
7.4	Chromatogram of residue from the interior of Boğazköy sherd 16.	183
7.5	The abundances of <i>n</i> -alkanes, alcohols and wax esters in six archaeological samples and the relative abundances of <i>n</i> -alkanes and wax esters in modern raw beeswax.	184

7.6	Chromatogram of the residues from the exterior of sherd 29 from Boğazköy.	185
7.7	The abundances of the main beeswax ingredients yielded by the interior and exterior surfaces of sherds 37 and 29 from Boğazköy.	187
7.8	The abundances of the main beeswax ingredients yielded by internal and external surfaces of sherds 21 and 24 from Boğazköy.	190
7.9	The structures of phytane, pristane, 17 α ,21 β -30-norhopane and the sterane 5 α -cholestane.	194
7.10	The chromatogram of the residue from the interior of sherd 3 from Hala Sultan Tekke, Cyprus.	195
7.11	The partial ion chromatogram for m/z 71 for the residue from the interior of Hala Sultan Tekke sherd 3.	196
7.12	The partial ion chromatogram for m/z 191 for the residue from the interior of sherd 3 from Hala Sultan Tekke.	197
7.13	The partial ion chromatogram for m/z 217 from the interior residue yielded by sherd Hala Sultan Tekke sherd 3.	198
7.14	(a) Total ion chromatogram for the non-saponifiable fraction of the residue from the interior of sherd 6 from Kouklia, (b) Partial ion chromatogram for m/z 71 and (c) Partial ion chromatogram for m/z 191 for the same residue.	200
7.15	Mass spectrum and structure of the trimethylsilyl (TMS) derivative of C _{16:0} fatty acid.	206
7.16	Mass spectrum and structure of the TMS derivative of oleic acid, one of the common isomers of C _{18:1} .	207
7.17	Fatty acid distributions in four sherds – Kuşakli sherd 1, Boğazköy sherd 28, Kalavassos sherd 8 and Kazaphani sherd 5 – where C _{8:0} and C _{9:0} fatty acids are present.	209
7.18	Mass spectrum and structure of the TMS derivative of azeleic acid (C _{9:0} dicarboxylic fatty acid).	210
7.19	The chromatogram produced by the residue extracted from the interior of sherd 8 from Kalavassos, Cyprus.	211
7.20	Mass spectrum and structure of the tris-TMS derivative of 9,10-dihydroxyoctadecanoic acid.	212
7.21	Mass spectrum and structure of the tris-TMS derivative of 11,12-dihydroxyoctadecanoic acid.	212
7.22	Mass spectrum and structure of the bis-TMS derivative of C _{18:0} 1-monoacylglycerol.	214
7.23	Mass spectrum of the bis-TMS derivative of C _{18:1} , 1-monoacylglycerol identified in the residue from the interior of Arpera sherd 1.	215

7.24	Mass spectrum of the bis-TMS derivative of C18:0 2-monoacylglycerol identified in the residue from the interior of Arpera sherd 4.	215
7.25	Mass spectrum and structure of the bis-TMS derivative of C16:0 1-monoacylglycerol showing the fragmentation producing the base peak at m/z 371.	216
7.26	Mass spectrum and structure of stigmasta-4,6,22-trien-3 β -ol identified in the residue from the interior of sherds 205 and 206 from Kilise Tepe, Turkey.	219
7.27	Chromatogram produced by the residue from the exterior of sherd 6 from Hala Sultan Tekke, Cyprus.	231
7.28	Mass spectra and structures of the TMS derivatives of (a) dehydroabietic acid, (b) pimaric acid, (c) sandaracopimaric acid and (d) 7-oxo-dehydroabietic acid.	232
7.29	Mass spectrum and structure of the TMS derivative of isopimaric acid identified in the residue from the interior of Kuşaklı sherd 4.	233
7.30	Chromatogram of the residue from the interior of sherd 1 from Hala Sultan Tekke, Cyprus.	236
7.31	Mass spectrum of (a) diethyl phthalate, (b) dibutyl phthalate and (c) dioctyl phthalate. Structures of (d) a generic phthalate molecule and (e) the fragment formed by re-arrangement in the mass spectrometer which produces the distinctive mass fragment at m/z 149.	237
7.32	Chromatogram produced by the residue from the interior of Kazaphani sherd 4 (a) and the mass spectra and structures of DDT(b), DDD(c) and DDE(d).	238
7.33	Chromatogram produced by the residue from the interior of sherd 1 from Enkomi, Cyprus.	240
7.34	Mass spectra and structures of (a) squalene and (b) the TMS ether of cholesterol.	244
7.35	Chromatograms of the TMS derivatives of visible residues from RLWm ware spindle bottles in (a) the Royal Ontario Museum, Canada, and (b) the Medelhavsmuseet, Stockholm, Sweden.	254
7.36	The total ion chromatograms of the TMS derivatives of the two non-saponifiable fractions prepared from the residue from (a) the Royal Ontario Museum, Toronto, Canada and (b) the Medelhavsmuseet, Stockholm, Sweden. (c) shows the partial ion chromatogram for m/z 71 extracted from chromatogram (b).	256
7.37	Chromatograms of the TMS derivatives of the saponified fractions of (a) The Royal Ontario Museum residue and (b) the Medelhavsmuseet residue.	257

7.38	Region of the chromatograms of the saponified fractions from (a) Royal Ontario Museum residue and (b) Medelhavsmuseet residue between 22 and 26 minutes retention time showing a range of hydroxy fatty acids.	258
7.39	Mass spectra and structures of (a) the bis-TMS methyl ester of 9,10-dihydroxyoctadecanoic acid, (b) the bis-TMS methyl ester of 11,12-dihydroxyoctadecanoic acid and (c) the tris-TMS derivative of 11,12-dihydroxyeicosanoic acid.	259
7.40	The mass spectra of (a) the peak eluting at 21.97 minutes in the chromatogram of the residue from the Medelhavsmuseet, (b) bis-TMS ricinoleic acid and (c) the TMS derivative of C _{19:0} . The structure of ricinoleic acid is also shown with the fragmentation producing the prominent peak at <i>m/z</i> 187 in the mass spectrum.	260
7.41	The FT-IR absorption spectrum for the residue from Saqqara sherd 1.	266
7.42	The results of the GC-C-IRMS analysis of nine absorbed and one visible archaeological residues	270
7.43	Results of GC-C-IRMS analysis of archaeological residues with the results for modern oils and beeswax.	273
7.44	Plot of $\Delta^{13}\text{C}$ against $\delta^{13}\text{C}_{16:0}$ for the archaeological residues with the results for modern oils and beeswax.	276
7.45	The numbers of sherds yielding each residue, those with mixed residues, the total number of sherds yielding residues and those which yielded no residues.	302
7.46	The total numbers of sherds yielding each of the four types of residues and no residues.	302
7.47	The number of residues of each type yielded by each vessel form.	309
7.48	The percentage of each residue yielded by each vessel form.	309
7.49	The number of RLWm ware sherds from each site yielding each of the four residues and no residues.	310
7.50	(a) The percentage of sherds of each form present in the combined assemblage containing material from this and a previous project and (b) the number of sherds in the combined assemblage yielding each of the four residues and no residues.	312
7.51	The percentage of each residue yielded by each vessel form for the combined assemblage.	313
7.52	The number of RLWm ware sherds from each site yielding each residue and no residue for the combined assemblage.	314
7.53	Results of compound specific stable isotope results for modern oils and modern raw beeswax.	318
7.54	Graph showing the results for $\Delta^{13}\text{C}$ against $\delta^{13}\text{C}_{16:0}$ for modern oils and modern raw beeswax.	321

7.55	Chromatogram produced by the TMS derivative of beeswax 1.	323
7.56	Chromatogram produced by the TMS derivative of beeswax 2.	323
7.57	Chromatogram produced by the TMS derivative of (a) a control sample of olive oil, (b) coriander oil after one steeping and (c) coriander oil after three steepings.	326
7.58	Analysis of frankincense oil.	328
7.59	A principle component analysis of the NAA data for 74 RLWm ware sherds and 11 local sherds.	331
7.60	Principle component analysis of the NAA data for 74 RLWm ware sherds and 11 local sherds.	333
7.61	Scatter plots showing the variation of (a) rubidium vs chromium and (b) scandium vs chromium for the 74 RLWm ware and 11 local samples.	335
7.62	Principle component analysis of LA-ICP-MS data for 5 RLWm ware sherds and 6 test sherds.	337
7.63	Second principle component analysis of the LA-ICP-MS data for 5 RLWm ware sherds and 6 test sherds.	338
7.64	Photographs of thin sections showing the presence of the self slip.	340
7.65	Photographs of thin sections showing textural features.	341
7.66	From Schubert and Kozal (2007), figure 2 – a principle component analysis of NAA results.	343
7.67	From Shubert and Kozal (2007), figure 3 – a scatterplot of scandium vs chromium from the NAA results.	344
A1.1	Chrononological chart, after Eriksson (1993, fig. 2, 4)	361
A1.2	Eriksson's revised chronology for LCIA to LCIIIB after Eriksson (2007, figure, 58)	362
A1.3	Relative chronologies for the LBA Aegean, after Warren and Hankey (1989), p71, Table 2.6).	363
A1.4	Absolute and relative chronologies for the Aegean, after Warren and Hankey (1989, 169, table 3.1)	364
A1.5	Relative and absolute chronologies for the Aegean Bronze Age after Shelmerdine (2008, fig. 1.1, 4)	365
A1.6:	A summary of the Low and High chronologies for the LBA Aegean after Shelmerdine (2008, fig. 1.2, 4)	366
A1.7	A comparison of chronologies for the LBA in the eastern Mediterranean	367